

A closer look at aortic seat belt injuries: review of 52 cases published in the last 60 years

Eduardo Silva , Vânia Constâncio , Celso Nunes , Leonor Baldaia , Miguel Castro , Luís Orelhas ,
Mário Moreira , Manuel Fonseca

Department of Angiology and Vascular Surgery, Coimbra Hospital and University Center; Coimbra, Portugal.

Submitted: May 7, 2023; Reviewed: May 30, 2023; Accepted: July 13, 2023

Presented at: 22nd Annual Meeting of the Portuguese Society of Angiology and Vascular Surgery 2023; Coimbra, Portugal.

ABSTRACT

INTRODUCTION: Seat belt aorta is a rare but often severe complication arising from blunt trauma with compression of the abdominal aorta against vertebrae. Seat belt sign is often present as an ecchymosis across the abdomen. The association with abdominal wall disruption and hollow viscus injury has been named seat belt triad; the presence of *Chance* fracture is sometimes considered a fourth component.

METHODS: Using Pubmed and Embase databases we reviewed all articles regarding abdominal seat belt aortic injuries and analysed presentation at admission, concomitant lesions, including presence of seat belt triad, treatment and outcomes.

RESULTS: Fifty-two cases were reported, from 1968 to 2019. Twenty-nine males (56%), mean age 43 ± 19 years. Most patients were stable at admission, with 29 (55.8%) presenting acute abdomen, 26 (50%) limb ischemia, 9 (17.3%) hypovolemic shock and 2 (3.8%) late-onset claudication. Seat belt sign was identified in 40 patients (76.9%), seat belt triad in 38 (73.1%) and 22 (42.3%) had *Chance* fractures, of which only 2 were not associated with seat belt triad. Most patients presented with aortic dissection (90.4%), complicated with pseudoaneurysm (11.5%), contained rupture (7.7%) or uncontained rupture (3.8%); 2 patients presented isolated iliac thrombosis and 3 limb ischemia. All patients required immediate surgical intervention, of which 40 (76.9%) required urgent vascular surgery. Forty-eight patients (92.3%) underwent vascular surgery: 39 open revascularizations and 9 endovascular procedures; three were managed conservatively. Ten patients (19.2%) passed away or died, of which 7 had seat belt triad. No patients needed reinterventions for vascular lesions except one, yet 3 required limb amputation. Most patients with seat belt triad required further visceral and abdominal wall repair.

CONCLUSIONS: Seat belt aorta and especially seat belt triad are severe complications associated with high morbimortality often requiring surgery and multiple interventions. As patients are usually conscious and stable upon admission, this condition should not be disregarded.

Keywords: Trauma; Seat Belt Triad; Seat Belt Aorta; Seat Belt.



INTRODUCTION

Seat Belt Aorta is a rare but potentially life-threatening injury that can occur in high-speed motor vehicle collisions. It refers to blunt trauma of the abdominal aorta, as a result of sudden deceleration caused by the seat belt during a collision, which compresses this large vessel against vertebrae. One of the key features of significant seat belt injuries is the presence of "Seat Belt Signs", which are visible signs of the injury. These marks appear as abrasions or bruises across the chest and abdomen and are caused by the force of the seat belt restraining the body during the collision, being indicative of internal lesions in up to 30% of cases (Figure 1).^[1]


Seat Belt Aorta is often associated with two other injuries collectively referred to as the "Seat Belt Triad", which are abdominal wall disruption and hollow viscus injury (Figure 2). A fourth component, more common in children, is sometimes present, which is a horizontal fracture of the vertebral body, due to flexion-distraction forces acting on the spine, called *Chance* fracture.^[2]

We aimed to review the published literature regarding aortic seat belt injuries.

Figure 1. Lower abdomen abrasions indicative of Seat Belt Sign.



Figure 2. Computed Tomography Angiography showing a patient with Seat Belt Triad



METHODS

Using *Pubmed* database we performed a non-systematic literature review on this subject including all the published cases reporting seat belt trauma to the abdominal aorta. We used different combinations of the search terms "seat belt aorta", "seat belt triad" and ("seat belt" AND "aorta"). All articles reporting blunt injury to the abdominal aorta due to seat belt trauma were included. We further analysed presentation at admission, concomitant lesions, including presence of seat belt triad and "*Chance* fracture", treatment and outcomes. Exclusion criteria were: articles not available in English or Portuguese; duplicated articles; reports on seat belt trauma to the thoracic aorta, the aortic arch or supra-aortic trunks; reports on paediatric population; blunt trauma to the aorta not resulting from seat belt use.

RESULTS

A total of 30 articles, published between 1968 and 2019 were included, of which only 12 (40%) were published after 2000. Fifty-two cases of seat belt injury to the abdominal aorta were reported, all of which were either case reports or case series.^[1-30] Twenty-nine males (56%) and 23 females (44%), with a mean age at presentation of 43 (SD ± 19) years, ranged 18 to 82.

Presentation at Admission and Clinical Features

All patients except one (n=51) presented at the emergency department in the acute phase, with the following symptoms: 29 (55.8%) with abdominal pain, 26 (50%) with acute lower limb ischemia and 9 (17.3%) with hypotension/hypovolemic shock. Two patients developed late-onset lower limb claudication (5 to 48 months after the accident).

The presence of concomitant abdominal wall and hollow viscus injury (seat belt triad) was reported in 38 patients (73.1%) and the presence of seat belt sign was identified in 40 patients (76.2%) at admission. Among these, only 7 patients with seat belt sign did not have seat belt triad ($p < 0.01$). Lumbar and *Chance* fractures were reported in 22 patients (42.3%), only 2 of which were not associated with the presence of seat belt triad ($p < 0.001$).

Twenty-four studies reported on 35 patients,^[1-21] 14 hemodynamic and conscience status at admission^[18-22,25-30] with 26 of these (74.3%) being hemodynamically stable when first entering the emergency department, while the remaining 9 (25.7%) were unstable. All patients were conscient at admission, except for one of the unstable patients, that was intubated and ventilated prior to arrival at the hospital.^[3]

The most frequent vascular lesion reported was intimal dissection of the infrarenal aorta and iliac arteries, present in 47 patients (90.4%), of which only 14 (29.8%) were uncomplicated dissections and the remaining 33 (70.2%) were complicated with either aorto-iliac thrombosis, false aneurysm formation, contained aortic rupture (peri-aortic hematoma), uncontained rupture or lower limb atheroembolism (Table 1).

Table 1. Reported vascular lesions caused by seat belt aortic injury, in 52 cases included in this review

Reported Vascular Lesions	Patients N=52
Aortic Dissection (All) – N (%)	47 (90.4)
Complicated Aortic Dissection – N (%)	33 (63.5)
Aortic Occlusion/Thrombosis	17 (32.7)
False Aneurysm	6 (11.5)
Contained Aortic Rupture (Periaortic hematoma)	4 (7.7)
Lower Limb Atheroembolism	4 (7.7)
Uncontained Aortic Rupture	2 (3.8)
Uncomplicated Aortic Dissection – N (%)	14 (26.9)
Isolated Lower Limb Atheroembolism – N (%)	3 (5.8)
Acute Iliac Thrombosis – N (%)	2 (3.8)
Other Reported Vascular Lesions – N (%)	6 (11.5)
Inferior Vena Cava Laceration	2 (3.8)
Superior Mesenteric Vein Tear	2 (3.8)
Iliac Vein Thrombosis	1 (1.9)
Inferior Mesenteric Artery Avulsion	1 (1.9)

Treatment and Outcomes

All patients presenting at the emergency department in the acute setting required immediate surgical intervention of some sort, with 40 (76.9%) requiring urgent vascular surgery and the remainder requiring either gastrointestinal, orthopaedic, abdominal wall repair or a combination of these. Eleven (21.2%) vascular lesions were initially managed conservatively,^[2,4,5,8,9,13,21,23,26] yet 3 required urgent surgery (1 to 3 days after admission), due to a rapidly expanding false aneurysm (n=2)^[8,21] or acute iliac thrombosis with limb ischemia (n=1)^[5] and 5 required later vascular repair (4 months to 2 years after admission) due to late-onset claudication (n=2),^[13,23] false aneurysm expansion (n=2)^[4,9] or acute iliac thrombosis with limb ischemia (n=1).^[26] One patient (1.9%) was too unstable and was considered for palliation only.^[17]

A total of 48 patients (92.3%) underwent vascular procedures, of which 39 (81.3%) open surgery and 9 (18.8%) endovascular repair (Table 2). The most common procedure was aortic graft interposition (n=12; 25%), followed by aortoiliac thromboendarterectomy (n=10; 20.8%), endovascular aortic repair (EVAR) (n=7; 14.6%) and aorto-femoral or aorto-iliac bypass (n=6; 12.5%). Forty-one patients (85.4%) required aortoiliac repair (32 open and 9 endovascular) and the remaining 7 (14.6%) underwent either extra-anatomical revascularization or lower limb procedures.

Table 2. Reported vascular operations performed in patients with seat belt aortic injury, of 48 cases included in this review

Procedure	Patients N=48
Aortic Graft Interposition – N (%)	12 (25)
Aortoiliac Thromboendarterectomy – N (%)	10 (20.8)
Endovascular Aortic Repair (EVAR) – N (%)	7 (14.6)
Aorto-femoral and Aorto-iliac Bypass – N (%)	6 (12.5)
Lower Limb Thromboembolectomy – N (%)	4 (8.3)*
Axillofemoral Bypass – N (%)	3 (6.25)
Palmar Aortic Stenting – N (%)	2 (4.2)
Direct Aortic Suture – N (%)	1 (2.8)
Not Indicated – N (%)	3 (6.25)

* Five additional lower limb thromboembolectomy were performed during other procedures.

Most patients with seat belt triad required multiple interventions for gastrointestinal and abdominal wall reconstruction, yet only 1 patient required vascular reintervention, due to a late-onset claudication 3 years after the initial surgery due to stent stenosis, which was treated by stent removal and interposition graft repair.^[13] Three patients required lower limb amputation, two of which were bilateral amputations.^[5]

Ten patients (19.2%) passed, on average 7.6 days after initial surgery (1 to 31 days), with a mean age of 61 years (36 to 78) and the majority (70%) being female (7:3). Out of these patients, 7 presented with seat belt triad at admission. Associated vascular lesions were: dissection complicated with acute aortic thrombosis (n=6); dissection complicated with uncontained aortic rupture (n=1) and uncomplicated aortic dissection (n=3). One patient died during surgery.

DISCUSSION

The increased use of computed tomography in adult trauma patients has led to a rise in the identification of blunt traumatic abdominal wall injuries. However, the incidence of blunt abdominal aortic injury is much lower, occurring in only 0.03% to 0.07% of blunt traumas due to the abdominal aorta's greater protection. Conversely, blunt aortic injury is more common, affecting around 0.58% of blunt trauma cases, with the majority of these injuries occurring in the thoracic aorta (>95%) and resulting in aortic rupture and death on site. Notably, seat belt use accounts for only a third of all cases of blunt abdominal aortic injury.^[5,13]

As it would be expected the most common injury to the aorta after blunt trauma was intimal dissection, which was the precursor in most cases of more severe complications such as false aneurysm formation, acute aortic thrombosis, and aortic rupture. This phenomenon, often results from sudden deceleration of the body during a high-speed accident causing the aorta to twist or stretch leading to mechanical stress on the aortic wall. Additionally, the seat belt can cause a sudden increase in intra-abdominal pressure, which can cause the aorta to expand and contribute to its dissection and rupture.^[4,13]

Despite most patients arriving at the emergency department hemodynamically stable and conscient, the majority required some sort of urgent or emergent surgical intervention. The presence of hemodynamically instability at admission was related to more severe vascular lesions, with 5 out of the 9 patients that presented with hypovolemic shock, having complicated aortic dissections (3 with a ruptured aorta and another 2 with pseudoaneurysm formation).

At the opposite end of the spectrum, clinically stable and asymptomatic aortic lesions might be tempting to manage conservatively, yet it should be taken into consideration possible complications arising from blunt aortic trauma, namely intimal dissection and false aneurysm formation. In fact, among the 11 patients managed conservatively only 3 did not need any kind of vascular surgical intervention, 2 of which had a small intimal flap aortic dissection and 1 had a partial iliac thrombosis.

Regarding the treatment, despite a clear preference towards open surgery, one should take into consideration that most of the presented cases were reported in the last century, with only 11 reported cases in the last 10 years, which is likely a reflection of progressive improvements made in automotive safety and legislation on road and traffic security, although one cannot exclude that this may also be due to publication bias. Interestingly, if we analyse the procedures by year, only 2 aortoiliac thromboendarterectomies and 4 aortic graft interpositions were performed after 2000 versus 6 EVAR. Independently of the procedure, almost all patients had good results regarding their initial vascular lesions, with only 1 patient requiring reintervention.

Lastly, one should take into account the high morbidity and mortality associated with seat belt aortic injuries and especially seat belt triad, with almost 20% of 30-day mortality reported in our study. These patients were generally older (mean age 61 versus 43 years) and female, yet male death was associated with younger patients (54 years, male mean age versus 64 years, female mean age), which might be due to men being more frequently involved in more violent and higher-speed vehicle accidents.

CONCLUSION

Abdominal aortic injuries resulting from seat belt trauma, including the seat belt aorta and the seat belt triad, are serious medical conditions that can have life-threatening consequences, being associated with a high morbidity and mortality. Despite patients often presenting stable

at admission, these injuries should not be disregarded. They require prompt recognition and management, often including multiple surgical interventions. As such, healthcare professionals should maintain a high degree of suspicion for these injuries and thoroughly evaluate patients who have been involved in high-speed motor vehicle accidents.

Acknowledgments: None

Conflicts of interest: None

Funding None

REFERENCES

- Cuesta RC, Casas IM, Armenteros FJ, Argote AX, Capitan JM. Seat Belt Syndrome and the Submarine Effect: A Case Report. *Chirurgia (Bucur)*. 2017;112: 624-6
- Kulvatunyou N, Albrecht RM, Bender JS, Friese RS, Bellal J, Latafi R, et al. Seatbelt triad: severe abdominal wall disruption, hollow viscus injury, and major vascular injury. *Am Surg*. 2011;77: 534-8
- Ahmed M, Mahmoud A, Samotowka M, Mitchell K, Saeed R. Seat Belt Aortic Dissection: A Case Report. *Cureus*. 2019;11:e4662
- Randhawa MP Jr, Menzoian JO. Seat belt aorta. *Ann Vasc Surg*. 1990;4:370-7.
- Freni L, Barbetta I, Mazzaccaro D, Settembrini AM, Dallatana R, Tassinari L, et al. Seat belt injuries of the abdominal aorta in adults--case report and literature review. *Vasc Endovascular Surg*. 2013;47:138-47
- Tomic I, Dragas M, Vasin D, Loncar Z, Fatic N, Davidovic L. Seat-Belt Abdominal Aortic Injury-Treatment Modalities. *Ann Vasc Surg*. 2018;53:270e16
- Warrian RK, Shoenuit JP, Iannicello CM, Sharma GP, Trenholm BC. Seatbelt injury to the abdominal aorta. *J Trauma*. 1988;28:1505-7
- Halancette M, Scalabrini B, Martinet O. Seat-belt aorta: a rare injury associated with blunt abdominal trauma. *Ann Vasc Surg*. 2006;20:681-3
- Campbell DK, Austin RF. Seat-belt injury: injury of the abdominal aorta. *Radiology*. 1969;92:123-4
- Clyne CA, Ashbrooke EA. Seat-belt aorta: isolated abdominal aortic injury following blunt trauma. *Br J Surg*. 1985;72:239
- Siriwardena AK. Seat-belt aortic injury. *Eur J Vasc Surg*. 1990;4:649-50.
- Ali MR Jr, Norcross ED, Brothers TE. Iliac and femoral artery occlusion by thromboemboli from an abdominal aortic aneurysm in the setting of blunt abdominal trauma. *J Vasc Surg*. 1998;27:545-8
- Gouveia E Melo R, Amorim P, Soares TR, Fernandes e Fernandes R, Ministro A, Garrido P, et al. Clinical aspects and present challenges of the seat belt aorta. *J Vasc Surg*. 2020;72:995-1004
- Shweiki E, Klena JW, Halm KA, Leonard DJ. Seat belt injury to the abdominal aorta. *Am J Emerg Med*. 2000;18:236-7
- Dunlop MG. Distal arterial emboli following seat belt injury of the aorta. *Injury*. 1986;17:370-1
- Johnstone BR, Waxman BP. Transverse disruption of the abdominal wall--a tell-tale sign of seat belt related hollow viscus injury. *Aust N Z J Surg*. 1987;57:455-60
- Frydenberg M, Royle JP, Hoare M. Blunt abdominal aortic trauma. *Aust N Z J Surg*. 1990;60:347-50
- Naude GP, Back M, Perry MO, Bongard FS. Blunt disruption of the abdominal aorta: report of a case and review of the literature. *J Vasc Surg*. 1997;25:931-5
- Gilani R, Saucedo-Crespo H, Scott BG, Tsai PI, Wall MJ Jr, Mattox KL. Endovascular therapy for overcoming challenges presented with blunt abdominal aortic injury. *Vasc Endovascular Surg*. 2012;46:329-31
- Thal ER, Perry MO, Crighton J. Traumatic abdominal aortic occlusion. *South Med J*. 1971;64:653-6
- Voellinger DC, Saddakni S, Melton SM, Wirthlin DJ, Jordan WD, Whitley D. Endovascular repair of a traumatic infrarenal aortic transection: a case report and review. *Vasc Surg*. 2001;35:385-9

- 22.** Reaney SM, Parker MS, Mirvis SE, Bundschuh CV, Luebbert PD, Vingan HL. Abdominal aortic injury associated with transverse lumbar spine fracture--imaging findings. *Clin Radiol.* 1995;50:834-8
- 23.** Danto LA, Wolfman EF Jr. Linear abdominal trauma. *J Trauma.* 1976;16(3):179-183.
- 24.** Bergqvist D, Takolander R. Aortic occlusion following blunt trauma of the abdomen. *J Trauma.* 1981;21:
- 25.** Crepps JT Jr, Rodriguez A. Combined abdominal aortic and visceral artery injury secondary to blunt trauma. *Injury.* 1989;20:319-22
- 26.** Vernhet H, Marty-Ané CH, Lesnik A, et al. Dissection of the abdominal aorta in blunt trauma: management by percutaneous stent placement. *Cardiovasc Intervent Radiol.* 1997;20:473-6
- 27.** Perry JR, Escobedo EM, Mann FA. Abdominal aortic injury associated with "seat belt syndrome". *Emerg Rad.* 2000;7:312-4
- 28.** Inaba K, Kirkpatrick AW, Finkelstein J, Murphy J, Brenneman FD, Boulanger BR, et al. Blunt abdominal aortic trauma in association with thoracolumbar spine fractures. *Injury.* 2001;32:201-7
- 29.** Stahlfeld KR, Mitchell J, Sherman H. Endovascular repair of blunt abdominal aortic injury: case report. *J Trauma.* 2004;57:638-41
- 30.** Sugimoto T, Omura A, Kitade T, Takahashi H, Koyama T, Kurisu S. An abdominal aortic rupture due to seatbelt blunt injury: report of a case. *Surg Today.* 2007;37:86-8