

Intestinal Ultrasound in Inflammatory Bowel Disease: A Valuable and Increasingly Important Tool

Catarina Frias-Gomes^a Joana Torres^{a, b} Carolina Palmela^a

^aGastroenterology Division, Surgical Department, Hospital Beatriz Ângelo, Loures, Portugal;

^bFaculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal

Keywords

Intestinal ultrasound · Diagnosis · Disease activity · Monitoring · Therapy response · Point-of-care

Abstract

Background: Intestinal ultrasound is emerging as a non-invasive tool for monitoring disease activity in inflammatory bowel disease patients due to its low cost, excellent safety profile, and availability. Herein, we comprehensively review the role of intestinal ultrasound in the management of these patients. **Summary:** Intestinal ultrasound has a good accuracy in the diagnosis of Crohn's disease, as well as in the assessment of disease activity, extent, and evaluating disease-related complications, namely strictures, fistulae, and abscesses. Even though not fully validated, several scores have been developed to assess disease activity using ultrasound. Importantly, intestinal ultrasound can also be used to assess response to treatment. Changes in ultrasonographic parameters are observed as early as 4 weeks after treatment initiation and persist during short- and long-term follow-up. Additionally, Crohn's disease patients with no ultrasound im-

provement seem to be at a higher risk of therapy intensification, need for steroids, hospitalisation, or even surgery. Similarly to Crohn's disease, intestinal ultrasound has a good performance in the diagnosis, activity, and disease extent assessment in ulcerative colitis patients. In fact, in patients with severe acute colitis, higher bowel wall thickness at admission is associated with the need for salvage therapy and the absence of a significant decrease in this parameter may predict the need for colectomy. Short-term data also evidence the role of intestinal ultrasound in evaluating therapy response, with ultrasound changes observed after 2 weeks of treatment and significant improvement after 12 weeks of follow-up in ulcerative colitis. **Key Messages:** Intestinal ultrasound is a valuable tool to assess disease activity and complications, and to monitor response to therapy. Even though longer prospective data are warranted, intestinal ultrasound may lead to a change in the paradigm of inflammatory bowel disease management as it can be used in a point-of-care setting, enabling earlier intervention if needed.

© 2021 Sociedade Portuguesa de Gastrenterologia
Published by S. Karger AG, Basel

Ecografia intestinal na doença inflamatória do intestino: uma ferramenta valiosa e de importância crescente

Palavras Chave

Ecografia intestinal · Diagnóstico · Atividade de doença · Monitorização · Resposta à terapêutica · Point-of-care

Resumo

Contexto: A ecografia intestinal na doença inflamatória intestinal tem ganho importância crescente como exame não invasivo para monitorizar a atividade de doença, pelos seus custos reduzidos, excelente perfil de segurança e disponibilidade. Neste artigo realizamos uma revisão sobre o papel da ecografia intestinal no manejo destes doentes.

Sumário: Na doença de Crohn, a ecografia intestinal tem uma boa acuidade no diagnóstico, avaliação da atividade e extensão da doença, assim como na avaliação de complicações, como estenoses, fístulas e abscessos. Apesar de não estarem validados, vários *scores* têm sido desenvolvidos para avaliar a atividade de doença. É de realçar a importância da ecografia intestinal na avaliação da resposta à terapêutica. A melhoria dos parâmetros ecográficos é observada tão precocemente como quatro semanas e persiste durante o seguimento a curto e longo prazo. Os doentes sem melhoria ecográfica parecem ter uma maior necessidade de intensificação terapêutica, corticóides, internamento ou cirurgia. À semelhança da doença de Crohn, a ecografia intestinal tem uma boa acuidade na avaliação ao diagnóstico, atividade e extensão da doença na colite ulcerosa. Na colite ulcerosa grave, um maior espessamento da parede intestinal à admissão está associado a maior necessidade de terapêutica de resgate e a ausência de melhoria deste parâmetro pode predizer a necessidade de colectomia. A ecografia também permite a avaliação da resposta à terapêutica na colite ulcerosa, com alterações observadas após duas semanas de tratamento e mantendo melhoria significativa após 12 semanas. **Mensagem-chave:** A ecografia intestinal é um método importante para avaliar a atividade de doença, complicações e monitorizar a resposta à terapêutica na doença inflamatória intestinal. Apesar de serem necessários mais estudos prospetivos, a ecografia intestinal pode levar a uma mudança de paradigma no manejo destes doentes, uma vez que pode ser utilizada no momento de prestação de cuidados, permitindo uma intervenção precoce quando necessário.

© 2021 Sociedade Portuguesa de Gastroenterologia
Published by S. Karger AG, Basel

Introduction

Objective evidence of bowel inflammation is a key feature in the management of inflammatory bowel disease (IBD) patients, since clinical-based assessment is insufficient to make adequate therapeutic decisions [1]. Endoscopic mucosal healing (MH) has emerged as a major therapeutic endpoint, as it has been associated with long-term clinical remission, steroid-free remission, and reduced risk of surgery [2, 3]. However, endoscopy is a time-consuming, expensive and invasive technique, not always tolerated by patients. Therefore, a growing interest has risen regarding non-invasive monitoring tools, such as intestinal ultrasound (IUS) and faecal calprotectin (FCal). IUS is a widely available imaging modality associated with low costs, an excellent safety profile, and lack of preparation [4]. It is increasingly recognised as an accurate technique as part of the armamentarium for IBD diagnosis, but also for assessing disease activity and extent, detecting complications, and monitoring response to therapy [5]. Moreover, IUS can be performed in a point-of-care setting, leading to therapy optimisation without delay, allowing repeated evaluations to monitor lesions over time, and even replacing invasive examinations, such as endoscopy [6]. Moreover, due to lack of radiation, good availability, and because it is an easy exam to perform for both patients and physicians, in experienced hands IUS can also replace other cross-sectional image modalities, such as computerised tomography (CT) or magnetic resonance (MR) [3]. When compared to other non-invasive monitoring tools such as FCal or C-reactive protein, IUS offers additional information, namely on disease extension, location, severity, and complications [4, 6]. Finally, in an era of shared decision-making with our patients, it is important to consider their acceptance when proposing follow-up examinations. In a recent systematic review, IBD patients preferred non-invasive techniques, particularly IUS, to monitor disease activity, when compared to endoscopy [7].

There is a clear need to consider IUS as a non-invasive monitoring tool in IBD, with recent ECCO-ESGAR recommendations supporting the use of IUS in the diagnosis and management of IBD patients [3]. In this review, we comprehensively discuss the role of IUS for: (a) screening and diagnosis of IBD; (b) evaluating disease activity and postoperative recurrence in Chron's disease (CD); (c) evaluating disease-related complications; and (d) monitoring response to therapy, both in CD and ulcerative colitis (UC) patients.




Fig. 1. Examples of IUS parameters. **a** Measurement of increased BWT (4.7 mm). **b** Increased CDF (Limberg score 4). **c** Areas of focal loss of BWS (asterisk). **d** Extramural findings, including mesenteric fat proliferation (arrows) and mesenteric lymph node (asterisk).

Screening and Diagnosis of IBD

IUS has been used as a screening tool in patients with gastrointestinal (GI) symptoms but without severe signs of disease (such as weight loss, anaemia, or elevated FCal), showing a good accuracy to distinguish IBD from irritable bowel syndrome patients in primary care settings [8]. Additionally, in a recent prospective study including 37 patients with low-risk abdominal symptoms, the use of IUS reduced the number of colonoscopies and appointments, improving health service outcomes [9]. Furthermore, GI infections can also mimic IBD. IUS has been shown to be an accurate method in the diagnosis of infectious enteritis when compared to CT or MR, and the major findings include hypoechoic small bowel wall thickening and lymph node enlargement. Similarly, IUS can also detect inflammation in infectious colitis. Importantly, all these IUS features may overlap with IBD, and IUS alone cannot diagnose GI infections. Therefore, an ultrasound control can be performed in these patients to exclude IBD [10].

The most frequent IUS parameter used to detect intestinal inflammation is bowel wall thickness (BWT; Fig. 1a). Common cut-off values are 2–3 mm for the small bowel and 3–4 mm for the colon [11]. Loss of bowel wall stratification (BWS) and increased vascularisation assessed through colour Doppler flow (CDF) are also associated with active inflammation (Fig. 1b, c) [11]. Finally, extramural features are also important, such as mesenteric fat proliferation and lymph nodes (Fig. 1d).

Therefore, IUS can be a very helpful tool for IBD diagnosis. For instance, CD patients at diagnosis should undergo small bowel assessment, either by MR enterography (MRE), IUS, and/or capsule endoscopy. CT enterography is another valid option, though it is associated with radiation exposure [3]. In a systematic review, including 1,558 CD patients, endoscopic, histologic, barium examination, and/or intraoperative findings were used for the reference standard. The overall pooled sensitivity of IUS was 88% and specificity 97% [12]. When specifically evaluating small bowel disease, the overall sensitivity of IUS ranged from 54 to 93%, with a specificity of 97–100% [13].

Table 1. Prospective studies evaluating IUS scores to predict disease activity or complications in CD

Index assessed (study)	n	Aims of the study	Segments assessed	Reference standard	Index parameters and cut-offs	Severity grades	Outcomes
UAICD (Futagami et al. [29], 1999)	55	To develop an IUS index of intestinal inflammatory activity	Duodenum Jejunum Ileum Ascending colon Transverse colon Descending colon Sigmoid colon Rectum	Colonoscopy Radiology	BWT >4 mm BWS Compressibility Peristalsis	A: Decreased compressibility and peristalsis with loss of haustration, but normal BWT B: BWT >4 mm and presence of BWS C: BWT >4 mm and loss of BWS	Strong correlation with endoscopy or barium contrast studies ($r = 0.62, p < 0.01$)
MS (Macconi et al. [35], 2003)	43	To establish whether IUS can assess histologic features of ileal stenosis in CD	NS	Surgical specimen Histology	Stenosis: BWT >4 mm with pre-stenotic dilation >25 mm BWS Sinus tract or fistulae	BW echo pattern: Hypoechoic pattern: increased BWT with loss BWS Stratified pattern: increased BWT with preserved BWS Mixed pattern: Co-existence of tracts with/without stratification	IUS detection of moderate-severe or intermediate degree of fibrosis: sensitivity 100%, specificity 63%
NYS (Neye et al. [31], 2004)	22	To evaluate diagnostic criteria of power Doppler sonography	Terminal ileum Caecum Ascending colon Transverse colon Descending colon Sigmoid colon	Ileocolonoscopy	BWT >5 mm CDF: no vessels/cm ² ; 1–2 vessels/cm ² ; >2 vessels/cm ²	Inactive: BWT <5 mm and no vessels Mild activity: BWT <5 mm and 1–2 vessels or BWT ≥5 mm with no vessels Moderate activity: BWT <5 mm and >2 vessels or BWT ≥5 mm with 1–2 vessels High activity: BWT ≥5 mm with 2 vessels	High concordance of power Doppler sonography and ileocolonoscopy (higher agreement in descending colon: $k = 0.91, 95\% \text{ CI } 0.83–0.98$)
PPRS (Paredes et al. [34], 2010)	33	To evaluate accuracy of IUS compared with endoscopy in the diagnosis and grading of postsurgical recurrence of CD	Anastomosis	Ileocolonoscopy	BWT (TI) >3 mm CDF: absent (0), barely visible (1), moderate vascularity (2), marked vascularity (3)	Recurrence: TI BWT >3 mm and/or positive CDF Moderate to severe recurrence: TI BWT >5 mm and/or CDF grade 2 or 3	Recurrence: sensitivity 76.9%, 95% CI 57.9–89; specificity 57.1%, 95% CI 25–84.2% Moderate to severe recurrence: sensitivity 86.7%, 95% CI 62.1–96.3; specificity 66.7%, 95% CI 43.7–83.7
LS LZS (Lenze et al. [36], 2012)	30	To compare FDG-PET/CT in stricture detection and stricture differentiation with IUS, endoscopy, and MRE	NS	Colonoscopy Histology	Stricture diagnosis: BWT >4 mm Bowel wall echogenicity CDF (Limberg score)	Stricture differentiation: Fibrotic: hyperechogenic BWT and Limberg 1 Mixed: mixed hypo and hyperechogenic BWT and Limberg 2 Inflammatory: hypoechogenic BWT and Limberg 3 or 4	Sensitivity of IUS to detect strictures: 68%, 95% CI 53–84 Correct diagnosis according to stricture differentiation: 40%
PCPRS (Paredes et al. [22], 2013)	60	To assess if CEUS can increase the value of IUS in the study of postoperative CD	Anastomosis	Ileocolonoscopy	BWT (TI) >3 mm CDF: absent (0), barely visible (1), moderate vascularity (2), marked vascularity (3) CEUS: % of wall brightness	0: normal BWT and CEUS enhancement <34.5% Recurrence: 1: BWT 3–5 mm with CEUS enhancement <46% 2: BWT >5 or CEUS enhancement >46% 3: BWT >5 mm or enhancement >70%, presence of fistulae	Strong correlation between IUS and endoscopy ($k = 0.64, p = 0.0001$) IUS sensitivity 89.8%, 95% CI 78.2–95.6; specificity 81.8%, 95% CI 52.3–94.9 CEUS sensitivity 98% (89.3–99.6), specificity 81.8% (52.3–94.5)

Table 1 (continued)

Index assessed (study)	n	Aims of the study	Segments assessed	Reference standard	Index parameters and cut-offs	Severity grades	Outcomes
US-LI (Rispo et al. [37], 2017)	71	To investigate the concordance between IUS-based Lémann Index (US-LI) and RME-based Lémann index (MR-LI)	Small bowel Colon Rectum	Ileocolonoscopy	BWT >3 mm BWS Stricture Abscesses and fistulae	Stricture: 1: BWT >3 mm or segmental enhancement without pre-stenotic dilation (SB; C) 2: BWT >4 mm or mural stratification without pre-stenotic dilation (SB; C) or <50% of the lumen (C) 3: BWT >4 mm, narrowed lumen and fluid distended (SB)/stricture with pre-stenotic dilation or >50% of the lumen (C) Penetrating disease: 2: Deep transmural ulceration 3: Hypochoic duct-like structures with fluid or air (SB)/phlegmon or any type of fistulae (C)	High concordance between US-LI and MR-LI ($r = 0.90, p < 0.001$)
SUS (Novak et al. [30], 2017)	63	To identify IUS parameters contributing to inflammatory disease activity, develop a simple score, and validate this score prospectively	Small bowel Terminal ileum Caecum Ascending colon Transverse colon Descending colon Sigmoid colon Rectum	Ileocolonoscopy	BWT (ileum >3 mm; colon >4 mm) CDF Mesenteric fat and lymph nodes Overall impression of disease activity	Continuous score: $[0.0563 \times \text{BWT1}] + [2.0047 \times \text{BWT2}] + [3.0881 \times \text{BWT3}] + [1.0204 \times \text{CDF1}] + [1.5460 \times \text{CDF2}]$	IUS overall sensitivity 92.1%, 95% CI 78.6–98.3, with 81.6% specificity, 95% CI 68–91.2 Score accuracy: AUC 0.836
RMS (Ramaswamy et al. [46], 2019)	35	To assess the utility of IUS in assessing disease activity in CD	Ileum Right colon Transverse colon Left colon Rectum	Ileocolonoscopy	BWT >3 mm CDF BWS Mesenteric fat Intestinal motility	Median BWT, Doppler activity and loss of BWS	Strong correlation with SES-CD ($r = 0.8, p = 0.009$)
IBUS-SAS (Novak et al. [33], 2021)	30	To establish the core parameters defining active intestinal inflammation in CD, to evaluate inter-rater reliability, and propose a segmental activity score	NS	Visual analogue scale	BWT CDF BWS Inflammatory fat	Continuous variable (ranging from 0 to 100) IBUS-SAS = $4 \times \text{BWT} + 15 \times \text{i-fat} + 7 \times \text{CDF} + 4 \times \text{BW}$	Inter-rater reliability: BWT 0.96, 95% CI 0.94–0.98 CDF 0.60, 95% CI 0.48–0.72 BWS 0.39, 95% CI 0.24–0.53 i-fat 0.51, 95% CI 0.34–0.67 IBUS-SAS 0.97, 95% CI 0.95–0.99
BUSS (Alloca et al. [32], 2021)	225	To assess the predictive value of bowel US findings and prospectively follow them up for a period of 12 months	Ileum Cecum-ascending colon Transverse colon Descending-sigmoid colon Rectum	Ileocolonoscopy	BWT CDF (0: absent; 1: present)	Continuous score (BUSS = $0.75 \times \text{BWT} + 1.64 \times \text{CDF}$) BUSS > 3.52 to predict endoscopic disease activity (AUC 0.864, 95% CI 0.812–0.906; sensitivity 83%, specificity 85%)	BUSS correlated significantly with SES-CD ($r = 0.55, p < 0.001$) BUSS > 3.52 predicted negative CD course (steroids, therapy optimisation, surgery, and hospitalisation)

BWT, bowel wall thickness; BWS, bowel wall stratification; CDF, colour Doppler flow; CEUS, contrast-enhanced ultrasound; MRE, magnetic resonance enterography; FDG-PET, fluoro-2-deoxy-D-glucose positron emission tomography; CT, computerised tomography; TI, terminal ileum; SB, Small bowel; C, colon; BW, bowel wall; NS, not specified.

Several studies have assessed the value of BWT to support the diagnosis of UC [4]. Even though UC is a mucosal disease, a BWT >4 mm had a sensitivity of 62–89% and specificity of 77–88% for its diagnosis [4]. Nevertheless, the best cut-off at diagnosis is not established and values >3 mm have also been reported.

In patients with active IBD, UC patients have a prominent thickening of the mucosal layer, whereas CD patients have a significant thickening of the submucosal layer and a higher rate of lymph node enlargement [14]. In UC, the thickening of the bowel wall is mostly proportional and BWS is usually present [11]. The mesenteric proliferation is a prominent feature in CD, although it can also occur in UC, especially during severe episodes [11]. Hence, IUS is an accurate method to screen for intestinal inflammation and to support the diagnosis of both CD and UC.

Evaluating Disease Activity in IBD

Disease Activity and Postoperative Recurrence in Crohn's Disease

IUS has shown a good accuracy in detecting disease activity in CD. In a systematic review, the overall sensitivity of IUS for assessing CD activity when compared to ileocolonoscopy, barium-contrasted exams, CT, MRE, capsule endoscopy, or surgical specimens was 89%, with a specificity of 94.3% [5], as previously reported [15]. When compared to MRE, IUS has an accuracy of 91% for localisation and 89% for bowel wall flow [16]. Similarly, in a recent prospective study, the accuracy of IUS was not significantly different from MRE, regarding BWT, loss of BWS and CDF, also highlighting the concordance between IUS and other cross-sectional exams [17]. The METRIC trial was a prospective multicentre trial including 284 patients (133 newly diagnosed; 151 relapsed) to evaluate MRE and IUS performance in assessing disease extent and activity in CD. A constructed referenced standard was used to compare the two techniques. Both MRE and IUS were highly accurate for detecting small bowel disease, even though a higher sensitivity and specificity in detecting disease activity and evaluating disease extent was observed with MRE [18]. Nonetheless, an expert panel highlighted some methodological limitations of this study such as bias in the constructed reference standard model, absence of information on time between MRE and IUS, and use of high BWT cut-offs [19]. Importantly, the sensitivity of IUS seems to be lower for jejunal lesions (55.6%) when compared to ileal (92.7%) or colonic involvement (81.8%) [5].

Regarding postoperative recurrence, even though ileocolonoscopy remains the gold standard examination, non-invasive tools may be considered, especially after small bowel resection [20]. In a recent systematic review, the pooled IUS sensitivity and specificity for detecting postoperative recurrence was 94 and 84% [21]. Small intestine contrast ultrasonography (SICUS) had a higher sensitivity (99 vs 82%), but lower specificity (74 vs. 88%) than IUS. Also, a higher concordance between contrast-enhanced ultrasound (CEUS) and colonoscopy has been observed when compared to IUS alone ($k = 0.82$ vs. 0.64 , $p < 0.001$), suggesting that both SICUS and CEUS can improve anastomosis evaluation [22]. Moreover, perianastomotic BWT correlated with Rutgeerts' endoscopic score ($r = 0.67$, $p = 0.0001$), with higher BWT in patients with a score ≥ 3 [23]. A cut-off BWT above 5.5 mm predicted severe endoscopic recurrence (≥ 3) [21].

Finally, a growing interest has emerged with the use of transperineal ultrasound (TPUS) to assess perianal disease as a simple and painless method. TPUS showed a sensitivity of 90.6% and a positive predictive value (PPV) of 93.4% in detecting perianal fistulae when compared to pelvic MR [24]. Extrasphincteric and suprasphincteric fistulae were less detected by TPUS, when compared to transsphincteric and rectovaginal/anovulvar fistulae. Regarding perianal abscesses, TPUS showed a sensitivity of 50% and PPV of 79% [25]. Importantly, although not completely studied, the steep learning curve of TPUS may limit the current use of this resource in clinical practice [26]. According to previous studies, physicians may achieve competency in TPUS after 12 months of training [27].

Accompanying the increasing evidence of IUS as an accurate tool to assess disease activity, several IUS scores have been published (Table 1). Six studies [28–33] evaluated inflammatory disease activity and showed a strong correlation between IUS score and endoscopy [28, 29, 31, 32]. Additionally, an expert consensus developed the International Bowel Ultrasound Segmental Activity Score (IBUS-SAS), with an almost perfect intraclass correlation coefficient (ICC 0.97 [0.95–0.99], $p < 0.001$) [33]. Nevertheless, the BWT definition varied between the studies, ranging from 3 mm [28], to 4 mm [29], or even 5 mm [31] in the colon. Additionally, two studies evaluated postoperative recurrence [22, 34], two compared stricture detection and echo pattern between IUS and MRE or histology [35, 36], and one investigated the concordance between IUS and MRE scores based on the Lémann index (LI) [37]. Interestingly, a high concordance was found between US-LI and MR-LI ($r = 0.90$, $p < 0.001$), suggesting

Table 2. Prospective studies evaluating IUS scores to predict disease activity in UC

Index assessed (study)	n	Aims of the study	Segments assessed	Reference standard	Index parameters and cut-offs	Severity grades	Outcomes
Ultrasound activity index (UAI) (Arienti et al. [65], 1996)	57 (severe or moderately severe pts)	To investigate IUS assessment of disease activity and extent	Ascending colon Transverse colon Descending colon Rectosigmoid colon	Tc-99m scintigraphy Surgical specimen	BWT	Continuous scale (sum of maximum BWT in four segments of the colon)	Strong correlation between Tc-99m scintigraphy and IUS ($r = 0.78, p < 0.001$) UAI: sensitivity 90.3%, specificity 96%
US score (Parente et al. [45], 2009)	74 (E1 pts excluded)	To evaluate colonoscopy and IUS as indexes of response to short-term therapy	Terminal ileum Colon	Colonoscopy	BWT >4 mm CDF	0: BWT <4 mm and no/sparse CDF 1: BWT 4-6 mm and CDF 2: BWT 6-8 mm and CDF 3: BWT >8 mm and CDF	Consistent concordance between endoscopy and IUS score in all visits (3, 6, and 9 months: k ranging from 0.76 to 0.90)
US score (Shikawa et al. [47], 2011)	37	To evaluate the association between sonoelastography (EG) and colonoscopy in assessing disease activity	Descending colon	Colonoscopy	BWT >4 mm BWS (presence or absence) EG (homogenous, random, hard)	Normal: BWT <4 mm Homogenous: BWT >4 mm, unclear BWS, homogenous EG Random: BWT >4 mm, unclear BWS, thickened wall EG with various colours Hard: BWT >4 mm, unclear BWS, homogenous blue EG	Significant association between EG and colonoscopy ($p < 0.001$)
US score (Civitelli et al. [43], 2014)	50 (paediatric E1 pts excluded)	To evaluate usefulness of IUS in assessing disease extent and activity	Right colon Transverse colon Left colon	Colonoscopy	BWT >3 mm CDF (presence or absence) BWS (yes or no) Absence of haustra coiled (yes or no)	Index calculation: sum of four components per segments 1: mild disease 2: moderate disease 3 or 4: severe disease	Strong correlation between IUS and endoscopy ($r = 0.94, p < 0.001$) US score >2 had a sensitivity of 100% and specificity of 93% to predict severe endoscopic disease
US score for UC (Hashimoto et al. [44], 2018)	116	Identify IUS parameters that can predict UC endoscopic activity and develop a simple US score	Ascending colon Transverse colon Descending colon Sigmoid colon	Colonoscopy	BWT (0: <3 mm; 10: 3-5 mm; 20: >5 mm) BWS (0 preserved, 2 obscure, 4 disappearing) CDF (Limberg score 0: 0; 1: 5; 2: 10; 3: 15)	Continuous score ranging from 0 to 39	UCUS showed a strong correlation with endoscopy (Mayo score: $r = 0.83, p < 0.001$; UCEIS score: $r = 0.85, p < 0.001$)
Hata index (Kinoshita et al. [48], 2018)	133	To evaluate IUS for assessing disease activity compared to colonoscopy	Caecum Ascending colon Right transverse colon Left transverse colon Descending colon Sigmoid colon	Colonoscopy	BWT BWS Ulceration	1: Normal BWT 2: Thickened mucosa and submucosa without hypoechoic changes of the submucosa 3: BWT with loss BWS 4: BWT with loss BWS and irregular mucosa or hypercholesteric shallow concavity in mucosa	IUS sensitivity 78.9%, specificity 63.8% Moderate concordance between IUS and colonoscopy ($k = 0.55, p < 0.001$) This study was posteriorly validated (Omotehara et al. [82])
Humanitas Ultrasound Criteria (HUC) (Alloca et al. [39], 2018)	53 (E1 pts excluded)	To assess diagnostic accuracy of IUS in detecting disease activity/severity and develop a non-invasive quantitative criteria of disease activity based on IUS findings	Ileum Ascending colon Transverse colon Descending colon Sigmoid colon	Colonoscopy	BWT >3 mm BWS (0 normal, 1 hypercholesteric, 2 loss) CDF (presence or absence) Lymph nodes Mesenteric hypertrophy	Index calculation: 1, 4 BWT + 2 x CDF Active disease (Mayo endoscopic score ≥ 2); HUC ≥ 6.3 points	BWT >3 mm and presence of CDF had a sensitivity of 68% and specificity of 100%, compared to colonoscopy HUC ≥ 6.3 points (sensitivity 71%, specificity 100%) to detect endoscopic active disease

Table 2 (continued)

Index assessed (study)	n	Aims of the study	Segments assessed	Reference standard	Index parameters and cut-offs	Severity grades	Outcomes
RM (Ramaswamy et al. [46], 2019)	102 colonic segments (number of patients not reported)	To develop a new IUS score in UC patients and assess its correlation with Mayo endoscopic score (MES)	Caecum Ascending colon Transverse colon Descending colon Sigmoid colon Rectum	Colonoscopy	BWT (0: <3 mm; 2: 3–5 mm; 4: >5 mm) BWT (0: present; 4: absence) CDF (0: no vessels; 2: 1–2 spots; 4: stretching in wall; 6: extending beyond the wall)	Continuous score Total score <4 = MCEIS 0/1 Total score 4–8 = MCEIS 2 Total score >8 = MCEIS 3	Excellent correlation between IUS and MES: caecum ($r = 0.95$), ascending colon ($r = 0.9$), transverse colon ($r = 0.96$), descending colon ($r = 0.85$), sigmoid colon ($r = 0.82$), rectum (0.76), $p < 0.001$
UC-IUS (Bots et al. [83], 2021)	60	To develop an ultrasound activity index	Ascending colon Transverse colon Descending colon Sigmoid colon	Colonoscopy	BWT >2 mm CDF (spots or stretch- es) Abnormal haustra- tions Fat wrapping	Continuous score ranging from 0 to 7 points	UC-IUS index showed a strong correlation with endoscopic Mayo score ($\rho = 0.83$, $p < 0.001$) and UCEIS index ($\rho = 0.759$, $p < 0.001$)

BWT, bowel wall thickness; BWS, bowel wall stratification; CDF, colour Doppler flow; CEUS, contrast-enhanced ultrasound; MCEIS, Mayo clinic endoscopic sub-score; Pts, patients; T, terminal ileum; UCEIS, ulcerative colitis endoscopic index of severity.

that IUS was not inferior to MRE to evaluate bowel damage.

Therefore, IUS is an accurate method to assess disease activity, even though a lower sensitivity when evaluating the jejunum has been shown. Regarding the postoperative setting, IUS is a useful method in detecting and grading the severity of recurrence in CD. Nevertheless, for patients with BWT <5.5 mm, IUS alone may not be sufficient to guide their management, as an accurate distinction between cicatrization and mild to moderate recurrence may not be achieved and, therefore, cannot replace endoscopy yet [38]. Finally, several endoscopic scores have been developed but none is fully validated. Accordingly, no specific IUS score is currently recommended to evaluate CD.

Disease Activity in UC

Although the role of IUS is less well established in UC, its value in evaluating disease activity has also been explored. In a prospective study, 53 UC patients underwent colonoscopy and IUS within 1 week. Patients with endoscopic active disease had higher BWT, presence of CDF, loss of BWS, and enlarged lymph nodes [39]. In a recent systematic review, most studies showed an association between IUS findings, either defined by BWT alone or in combination with other features, and disease severity on endoscopy [4]. Moreover, the accuracy of IUS to evaluate disease extension compared to endoscopy was reported as 88.5–95% (sensitivity 95%; specificity 96%) [4]. Assessments of the sigmoid and descending colon had the higher accuracy [40], in contrast to the rectum, where trans-abdominal IUS had a poor sensitivity (15%) [41]. Nevertheless, this limitation could be exceeded using TPUS. In a cross-sectional study, 57 UC patients underwent trans-abdominal and TPUS evaluation simultaneously, 7 days before or after colonoscopy. Rectal BWT ($r = 0.72$, $p < 0.001$) and CDF ($r = 0.66$, $p < 0.001$) correlated well with the Mayo endoscopic score, suggesting that TPUS can be a good tool to evaluate patients with proctitis [42].

Considering IUS scores, nine indices have been prospectively developed in UC (Table 2). All studies were based on BWT and usually complemented by CDF [39, 43–46] and/or BWS [39, 4, 44, 46, 48]. Most studies considered a normal BWT when below 3 mm, even though two studies considered 4 mm to define normal BWT [45, 47]. Overall sensitivity of UC scores ranged from 71 to 100% and specificity from 63.8 to 100%. A strong correlation was found between IUS scores and endoscopy [43–45, 47], especially in severe disease ($r = 0.94$, $p < 0.001$) [43].

Thus, IUS has shown a good performance in assessing disease activity in UC, although a lower sensitivity has been reported when evaluating the rectum, which could be exceeded using TPUS. Similar to CD, no IUS score has been formally validated.

Evaluating Disease-Related Complications

Crohn's Disease

Several studies have assessed IUS accuracy to detect intestinal strictures, with a sensitivity ranging from 74.4 to 100% and a specificity of 63–100% [35, 49–53]. Strictures have been defined by a thickening and stiffness of the bowel wall, accompanied by a proximal dilation >2.5 cm (Fig. 2) [54]. In a prospective study including 249 CD patients, the concordance between MRE and IUS for stricturing disease was high when compared to intraoperative findings ($k = 0.86$) [55]. SICUS seems to have higher sensitivity for detecting strictures when compared to IUS (89–94 vs. 74–76%) [56, 57] and showed a good accuracy in detecting ileal stenosis and prestenotic dilation [58, 59]. However, it is still not clear if IUS, including SICUS, can distinguish inflammatory from fibrotic stenosis. Nevertheless, assessment of the wall echo pattern at the stricture level may suggest the degree of fibrosis. Maconi et al. [35] concluded that strictures with a stratified echo pattern had a higher degree of fibrosis compared to those characterised by a hypoechoic echo pattern. Moreover, a reduced CDF has also been associated with a fibrotic phenotype [60]. Importantly, CEUS has also been reported as an adjuvant method to characterise strictures in CD. When compared to surgical specimens, the concordance between CEUS with inflammatory or fibrostenotic phenotype was good ($k = 0.63$), with a good correlation between sonographic and pathology scores for both inflammatory ($r = 0.53$, $p = 0.004$) and fibrotic stenosis ($r = 0.50$, $p = 0.007$) [60]. Finally, conflicting data have been published when evaluating sonoelastography as a possible method to distinguish fibrotic from inflammatory strictures in CD, and this modality requires further investigation [61].

Penetrating disease is another potential complication in CD. Abscess appears as an irregular hypoechoic lesion without vascularisation. Fistulae are hypoechoic tracts, originating from the bowel wall and connecting to other tissues, such as the urinary bladder, skin, vagina, or other intestinal segments (Fig. 3) [54]. In a meta-analysis, the pooled sensitivity and specificity of IUS in detecting fis-




Fig. 2. IUS showing an ileal stenosis, with thickened bowel wall with narrow lumen (asterisk) and prestenotic dilation (arrow).

tulae was 74 and 95% and in diagnosing abscesses was 84 and 93%, respectively [15]. Ripollés et al. [62] showed that CEUS was able to differentiate between phlegmon and abscess in 57 CD patients, showing a high concordance ($k = 0.972$) with CT, MR, percutaneous drainage, or surgery. Similar findings have been previously reported, highlighting the role of CEUS as a sensitive method for differential diagnosis between phlegmon and abscess (Fig. 4) [63].

Therefore, IUS is an accurate method to evaluate CD-related complications. SICUS can help evaluating patients with strictures. CEUS supports the differential diagnosis of an inflammatory mass and is a promising tool in differentiating inflammatory from fibrotic strictures.

Ulcerative Colitis

A particular important scenario is acute severe UC (ASUC), treated with high-dose systemic corticosteroids, which is associated with an increased risk of colectomy. Nowadays, therapy response is based on clinical symptoms and biochemical markers (Oxford criteria) [64]. In hospitalised patients with moderate to severe UC, a significant decrease in BWT was observed in all patients who did not require colectomy, whereas patients who underwent colectomy had no BWT improvement between admission until day 10 [65]. In a recent pilot study including 10 patients, higher BWT (6.2 vs. 4.6 mm, $p = 0.009$) and any colonic segment with BWT >6 mm at admission were also associated with the need for infliximab salvage therapy. Additionally, after 3 days of high-dose steroid therapy, steroid-responsive patients had lower BWT (4.0 vs. 6.3 mm, $p = 0.009$) [66]. Similarly, in a retrospective study including 69 ASUC episodes in 52 paediatric patients, sal-




Fig. 3. IUS showing entero-enteric fistulae: hypoechoic tracts connecting small bowel loops (arrows).




Fig. 4. Two examples of CEUS showing differentiation between abscess and inflammatory mass. **a** Using CEUS this hypoechoic mass shows three areas completely devoid of microbubble signal, repre-

senting three abscesses. CEUS can be very helpful for defining the size of the abscesses. **b** Using CEUS this hypoechoic structure shows intralesional enhancement and corresponds to an inflammatory mass.

vage therapy was more frequently needed in patients with higher BWT, higher vascularisation, and loss of BWS at admission. A thickened wall (>3.4 mm) and loss of BWS were independent predictors of steroid resistance [67]. Thus, if IUS parameters prove to be independent predictors of response to systemic steroids in the ASUC setting, early IUS could enable a timelier introduction of salvage therapy.

Monitoring Response to Therapy

Crohn's Disease

Several studies have assessed IUS as a monitoring tool in CD to evaluate the response to therapy [45, 68–70]. The definition of ultrasonographic remission, or transmural healing (TH), is not yet established, although it has been defined by some authors as a complete normalisation of

BWT (<3 mm) with normal CDF or a complete normalisation of all IUS parameters. Additionally, definitions for ultrasonographic response have also been proposed when a sonographic improvement occurs [71]. In a prospective study, TH was associated with higher rates of steroid-free remission, lower rates of clinical relapse, and longer intervals until hospitalisation when compared to MH, suggesting that TH may be a more accurate target than MH alone in CD [72]. In the recently published STRIDE-II update, TH is considered as a potential therapeutic target but not a formal one yet [1].

The TRUST study was a 12-month prospective study to assess the value of IUS in monitoring CD, including 243 patients with at least moderately active CD (Harvey Bradshaw index >7). A significant proportion of patients had an improvement in BWT, CDF, BWS, and mesenteric fat proliferation at the end of follow-up. These ultrasonographic changes were accompanied by

Table 3. Studies evaluating IUS as a monitoring tool to evaluate therapy response in CD and UC patients

Study	Participants and duration of follow-up	Study type	Primary endpoint	Inclusion criteria	IUS features	Response to treatment	Outcomes
<i>Crohn's disease</i>							
Ripollés et al. [68] (2016)	51 CD Median follow-up 16 months (IQR 12.2–32)	Multicentre prospective study	Assess long-term effect of biological treatment on transmural lesions by IUS (including CEUS)	Active CD pts with clinical indication for anti-TNF	BWT (>3 mm) CDF (0: absent; 1: barely visible; 2: moderate vascularity; 3: marked vascularity) Wall brightness after contrast enhancement	Clinical-biological response: Remission: HBI <5 and normal CRP levels, without steroids Partial response: HBI decrease >3 points and CRP levels decreased but without normalisation, without steroids Lack of response: HBI and/or CRP increased or did not change when other treatments were needed to control the disease IUS remission: BWT ≤3mm, CDF 0, no complications IUS improvement: BWT decrease ≥2 mm, CDF decrease 1 grade; decrease ≥20% enhancement, no complications	Week 12: Significant reduction in BWT in patients with clinical remission/partial response (4.66 vs. 1.79 mm, $p = 0.01$) Week 52: 96% of patients who showed a clinical remission or response had IUS improvement or normalisation Good IUS response at week 12 predicted good IUS response at week 52, with a sensitivity of 75.9% and specificity of 81.8% Follow-up: Patients without IUS improvement at week 52 had a higher need for therapy intensification or surgery (65 vs. 11%, $p < 0.001$)
Kucharzik et al. [70] (2017) TRUST study	234 CD 12 months	Multicentre prospective study	Change in IUS parameters within 12 months	CD pts who experienced a flare: HBI ≥7 points	BWT (TI >2 mm; colon >3 mm) CDF (Limberg score)	Clinical response: decrease in HBI score of 3 points Clinical remission: HBI <4	Changes in IUS parameters ($p < 0.001$) BWT: TI 75.4 vs. 35.8%, SC 47 vs. 23.1%; CDF (LS ≥3) 44 vs. 9.7%; BWS: 53% vs. 21.6%; mesenteric fibro-fatty proliferation 47 vs. 17.9% Improvement in IUS was accompanied by clinical (median HBI 10 vs. 2, $p < 0.001$) and biochemical response (median CRP 50.5 vs. 16.1 g/dL, $p < 0.001$)
Kucharzik et al. [73, 74] (2020) STARDUST IUS sub-study	82 CD week 16 71 CD week 48	Phase 3b randomised trial of CD patients treated with UST, comparing TZT vs. SoC	Changes in IUS parameters, including transmural response to UST	Moderate to severe active CD (CDAI 220–450), who failed conventional therapy and/or 1 biologic	BWT CDF BWS IF	IUS response: ≥25% BWT reduction from BL IUS remission: normal BWT, CDF, BWS, and absence of IF	IUS response: 33.8% (week 16); 35.8% (week 48) IUS remission: 11.3% (week 16); 18.3% (week 48) Mean BWT improvement from BL was observed as early as week 4 ($p = 0.0002$) BWT and CDF sign started to normalise at week 8; IF and BWS at week 16
Calabrese et al. [75] (2021)	188 CD	Multicentre prospective study	Assess changes in IUS parameters, including TH, induced by different biological therapies	Patients eligible for biological therapies	BWT (small bowel <3 mm; large bowel <4 mm) CDF BWS Disease length Lymph nodes Fibro-fatty proliferation	Improved lesions: improvement (>1 mm) or normal BWT; decreased length of disease; Limberg score improvement; no worsening of other parameters TH: normalisation of all parameters	3 months: improved lesions: 36%; TH 16.4% 6 months: improved lesions: 38%; TH 24.5% 12 months: improved lesions: 36%; TH: 27.6% Colonic lesions: higher risk of TH at 3 months (OR 3.18, 95% CI 1.16–7.75) Greater BWT: lower rates of TH at 3 and 12 months (OR 0.70, 95% CI 0.5–0.97; 0.58, 95% CI 0.38–0.89)

Table 3 (continued)

Study	Participants and duration of follow-up	Study type	Primary endpoint	Inclusion criteria	IUS features	Response to treatment	Outcomes
<i>Ulcerative colitis</i>							
Maconi et al. [69] (1999)	30 UC 2 months (E1 patients excluded)	Prospective study	Determine whether IUS evaluation of BWT may be useful in follow-up of UC	Active UC	BWT >4 mm Absence of regular haemorrhage	Remission: no symptoms and/or no signs of disease activity on endoscopy	BWT decreased in patients who achieved clinical remission (7.3 vs. 5.1 mm, $p < 0.001$) after treatment and did not change in those patients without significant clinical improvement (7.0 vs. 7.0, $p = NS$)
Parente et al. [45] (2009)	74 UC 15 months (E1 patients excluded)	Prospective study	Evaluate the accuracy of IUS as a surrogate of colonoscopy in monitoring response to medical therapy	Recently diagnosed or flare-up UC patients, with moderate-to-severe disease and needing high-dose systemic steroids (oral or IV)	BWT BWS CDF	Endoscopic remission: Bs: 0 Endoscopic relapse: Bs > 1 US severity: 0: BWT <4 mm and no/sparse CDF 1: BWT 4–6 mm and CDF 2: BWT 6–8 mm and CDF 3: BWT >8 mm and CDF	Consistent concordance between 0–1 Baron scores and US scores in the 3rd, 9th, and 15th months (k ranged from 0.76 to 0.90) Severe IUS scores (2–3) after 3 months of therapy had a higher risk for severe endoscopic activity in the 15th month (OR 9.1, 95% CI 2.5–33.5)
Maaser et al. [79] (2020) TRUST&UC study	224 UC 12 weeks	Multicentre prospective study	Proportion of patients with normalisation of BWT in patients with clinical response	UC patients in clinical relapse (SCCAI ≥ 5 points)	BWT (>3 mm, except in sigmoid colon >4 mm) CDF (present or absent)	Clinical response: decrease ≥ 3 points in SCCAI	Significant reduction at week 12 in BWT (SC [89 vs. 32%] and DC [83 vs. 37.6%]) and in CDF (SC [34.8 vs. 12.9%] and DC [15.2 vs. 7.3%]) Patients with a normalisation of BWT had higher rates of clinical response (SC: 90.5 vs. 68.9%; DC: 96.4 vs. 68.8%, $p < 0.001$)
Sacrallo et al. [67] (2020)	52 UC (paediatrics)	Retrospective	Evaluate the potential role of IUS in predicting the need for second-line therapy in ASUC	ASUC patients (PUCAI >65)	BWT BWS CDF Lymph nodes	Steroid treatment failure: need for second-line therapy (infliximab or calcineurin inhibitor)	Patients requiring a second-line therapy had higher BWT values (5.14 vs. 3.89 mm, $p < 0.001$) Loss of BWS was more frequent in steroid-resistant patients (47 vs. 3%, $p < 0.001$) BWT >3.4 mm predicted steroid treatment failure, with a sensitivity of 92% and specificity of 52%
Smith et al. [66] (2021)	10 UC	Prospective study	IUS can predict steroids-refractory disease	ASUC patients (>6 bowel wall movements/day)	BWT (>4 mm) CDF (Limberg score) BWS Extra-intestinal features	Steroid treatment failure: need for salvage therapy with infliximab	At admission, BWT was higher in patients with steroid treatment failure (6.2 vs. 4.6 mm, $p = 0.0009$). Patients with any colonic segment >6 mm were more likely to require salvage therapy (100 vs. 25%, $p = 0.03$) At day 3, steroid responsive group had lower BWT (4.0 vs. 6.3 mm, $p = 0.0009$)

BWT, bowel wall thickness; BWS, bowel wall stratification; CDF, colour Doppler flow; TH, transmural healing; CEUS, contrast-enhanced ultrasound; Pts, patients; TI, terminal ileum; SC, sigmoid colon; DC, descending colon; T21, treat to target; SoC, standard of care; UST, ustekinumab; CDAI, Crohn's disease activity index; IF, inflammatory fat; SCCAI, Short Clinical Colitis Activity Index; IV, intravenous; Bs, Baron score; ASUC, acute severe ulcerative colitis; BL, baseline.

clinical and biochemical improvement [70]. Similarly, in a multicentre prospective study, improvement of BWT and CDF were observed after 12 weeks, increasing even more after 12 months of therapy, highlighting that IUS response at week 12 was associated with maintenance of the IUS response at week 52 [68]. Importantly, patients without IUS improvement after 1 year of therapy had a higher need for therapy intensification or surgery (65 vs. 11%, $p < 0.001$) [68]. Likewise, in an interim analysis of the STARDUST trial IUS sub-study including 88 CD patients, IUS response and remission after ustekinumab induction were assessed. IUS response was defined by a BWT reduction of 25% from baseline and IUS remission by normalisation of BWT, CDF, BWS, and inflammatory mesenteric fat. At week 16, IUS response and remission rates were 33.8 and 11.3%, respectively. BWT improvement was observed as early as week 4, suggesting that IUS could be a useful method to detect early response to treatment [73]. A consistent decrease in BWT was observed up to week 48. Furthermore, the overall IUS response progressively increased over time (week 48 46.3%), accompanied by a higher rate of TH (week 48 24.1%). Interestingly, normalisation of BWT was more frequent when the colon was affected compared to the ileum (50 vs. 15.8% at week 48), reflecting a faster cicatrization of the colon [74]. A recent multicentre prospective study, including 181 CD patients treated with different types of biologic therapies, assessed IUS improvement (decrease ≥ 1 mm or normalisation of BWT, decrease in length of disease, Limberg score improvement, and no worsening of other IUS parameters) and TH (normalisation of all parameters) during 12 months of follow-up. After 3 and 12 months, 36.7 and 36% of the patients showed IUS improvement, with 16.4 and 27.6% achieving TH, respectively. Patients in clinical and biochemical remission had higher rates of TH. Predictive factors of TH included colonic location (aOR 3.18, 95% CI 1.11–9.10), whereas greater BWT at baseline was associated with lower rates of TH at 3 (aOR 0.70, 95% CI 0.5–0.97) and 12 months (aOR 0.58, 95% CI 0.38–0.89) [75]. Similarly, in a recent prospective study, baseline BWT and CDF, presence of disease-related complications, FCal (>250 $\mu\text{g/g}$), and male gender were associated with a higher need for steroids, optimisation therapy, hospitalisation, or surgery after 12-months of follow-up [31]. Thus, IUS features at baseline and IUS improvement during follow-up seem to be associated with disease-related outcomes. In a prospective study, including 80 consecutive CD patients, baseline and follow-up SICUS

were performed (after a median of 18 months). Patients with IUS response (improvement or normalisation of BWT, decreased length of disease, without complications) had lower need for steroids, hospitalisation, and/or surgeries at 1 and 5 years of follow-up [76]. Regarding CEUS, differences in kinetic parameters derived from time intensity curves, such as peak enhancement, wash-in perfusion index, wash-in and wash-out rate, significantly improved in patients with clinical or endoscopic response, after 6 weeks of therapy [77]. Similarly, in a prospective study of IBD patients treated with vedolizumab, amplitude-derived CEUS parameters of mural microvascularisation also decreased in clinical responders after 14 weeks of therapy [78]. Altogether, these data emphasise the role of IUS as a method for monitoring the response to treatment in CD patients (Table 3).

Ulcerative Colitis

In the TRUST&UC prospective study, IUS findings in UC patients after initiating therapy for clinical relapse were evaluated during a 12-week period [79]. Overall, 178 patients with left-sided or pancolitis completed follow-up at week 12. Patients with normalisation of BWT in the sigmoid or descending colon had higher rates of clinical response. Moreover, clinical responders showed a significant reduction in BWT and CDF at week 12. These changes could be observed as early as after 2 weeks of therapy [79]. Finally, other IUS parameters, such as mesenteric fat proliferation, BWS, haustration, and ascites also improved after 12 weeks. Clinical symptoms accompanied IUS improvement, with a lower Simple Clinical Colitis Activity Index (SCCAI) at week 12 (9 vs. 2 points, $p < 0.001$). Similarly, a higher proportion of patients with BWT normalisation at week 12 had normal FCal values (<250 $\mu\text{g/g}$; sigmoid colon: 48.9 vs. 22.2%, $p = 0.02$; descending colon: 50 vs. 25%, $p = 0.03$) [79]. Parente et al. [45] also evaluated moderate to severe UC patients during a 15-month follow-up period. Patients who had severe IUS activity in the third month after corticosteroids therapy had a higher risk of severe endoscopic activity at 15 months (OR 9.1, 95% CI 2.5–33.5; Table 3).

Even though studies with longer follow-up are needed, these data support the use of IUS as a non-invasive monitoring tool to assess therapy response in UC. Importantly, the IUS response can be observed as early as 2–4 weeks after treatment initiation.

Current role of IUS in IBD
Screening and diagnosis of IBD Differential diagnosis with IBS and GI infections Small bowel assessment in CD Disease activity assessment in IBD Evaluation and grading of disease activity and extent Severe postoperative recurrence in CD Perianal disease in CD-TPUS Disease-related complications in CD Strictures, fistula such as and inflammatory masses, such as phlegmon and abscesses Monitoring response to therapy in IBD
Future directions
Validation of IUS scores to allow its use in clinical practice and clinical trials Improve IUS assessment of mild to moderate CD postoperative recurrence CEUS as a promising tool differentiating fibrotic from inflammatory strictures Transmural healing as potential target in CD Uniform time points for IUS evaluation based on a treat-to-target strategy IUS in ASUC: predict need for salvage therapy and colectomy Correlation of IUS with histology in UC

Fig. 5. The current role and future directions of IUS in IBD. ASUC, acute severe ulcerative colitis; CD, Crohn's disease; GI, gastrointestinal; IBS, irritable bowel syndrome; TPUS, transperineal ultrasound; UC, ulcerative colitis.

Future Directions and Conclusions

Nowadays, IUS is a very useful tool in the management of IBD patients, with a good accuracy in detecting disease activity, extent, and complications in CD. Besides, although being a mucosal disease, recent published data also endorse its use in UC to assess disease activity and extension. Emerging data have supported the use of IUS as a promising tool to assess response to treatment in both UC and CD, reporting changes in IUS features as early as 2–4 weeks of treatment and that persist in short- and long-term follow-up (Fig. 5). In fact, this could lead to a paradigm change in IBD, as IUS can become a routinely used tool in the management of these patients in a point-of-care setting and enabling early intervention if needed. Nevertheless, the use of IUS is not yet universal and its performance is highly dependent in the operator's experience. Inter-observer agreement of IUS in UC and CD patients is excellent for BWT and good for CDF, with fair to moderate agreement in other IUS parameters, such as lymph nodes and inflammatory fat [80, 81]. Moreover, IUS can have lower accuracy in specific bowel locations, such as the proximal jejunum and rectum. Other possible limitations of IUS include the patient's biotype, as evaluation in obese patients is difficult [82], evaluation of disease activity/extent in the postoperative setting, due to anatomical changes, and the lower capacity to detect superficial lesions in the small bowel. Therefore, it is important to train IBD-specialised gastroenterologists in this technique, as proposed by the International Bowel Ultrasound (IBUS) group. Additionally, future studies are

needed to improve IUS capacity in differentiating the severity of endoscopic recurrence in the postoperative setting, as well as to deepen the knowledge on elastography and better characterisation of stricture subtype in CD. Regarding UC, the real accuracy of IUS to predict histologic remission has never been formally studied. In an era of strict endpoints like endoscopic Mayo score of zero or even histological remission, IUS parameters might not be sensitive enough to capture subtle inflammatory mucosal changes. Finally, no IUS score has been fully validated and a homogenous approach of IUS parameters is warranted to spread its use in IBD clinics and hospitals, as well as in clinical trials.

In conclusion, IUS is an accurate non-invasive monitoring tool not only to assess IBD diagnosis, disease extent, and activity in CD and UC, but also to monitor response to therapy. In experienced hands, IUS adds extraordinary value to the management of IBD patients.

Acknowledgements

We would like to thank Dr. Frauke Peterson from the University Teaching Hospital of Lueneburg, Germany, for kindly sharing her CEUS cases for this review paper.

Statement of Ethics

This article does not contain any studies with humans or animal subjects performed by the authors.

Conflict of Interest Statement

J.T. received speaker and consulting fees from Janssen and consulting fees from Arena Pharmaceuticals, and Gilead and Galapagos. C.P. received speaker and consulting fees from Janssen and Laboratórios Vitória.

Funding Sources

This review had no funding resources.

Author Contributions

All the authors made substantial contributions to the article. C.F.-G., J.T., and C.P. contributed to the manuscript concept and design. C.F.-G. drafted the article, and J.T. and C.P. made important revisions and contributions. All authors critically revised the manuscript. All authors have approved the final version of the manuscript.

References

- 1 Turner D, Ricciuto A, Lewis A, D'Amico F, Dhaliwal J, Griffiths AM, et al.; International Organization for the Study of IBD. STRIDE-II: An Update on the Selecting Therapeutic Targets in Inflammatory Bowel Disease (STRIDE) Initiative of the International Organization for the Study of IBD (IOIBD): Determining Therapeutic Goals for Treat-to-Target strategies in IBD. *Gastroenterology*. 2021 Apr;160(5):1570–83.
- 2 Shah SC, Colombel JF, Sands BE, Narula N. Mucosal healing is associated with improved long-term outcomes of patients with ulcerative colitis: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. 2016 Sep;14(9):1245–1255.e8.
- 3 Maaser C, Sturm A, Vavricka SR, Kucharzik T, Fiorino G, Annesse V, et al.; European Crohn's and Colitis Organisation [ECCO] and the European Society of Gastrointestinal and Abdominal Radiology [ESGAR]. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 1: initial diagnosis, monitoring of known IBD, detection of complications. *J Crohns Colitis*. 2019 Feb;13(2):144–64.
- 4 Smith RL, Taylor KM, Friedman AB, Gibson RN, Gibson PR. Systematic Review: Clinical Utility of Gastrointestinal Ultrasound in the Diagnosis, Assessment and Management of Patients With Ulcerative Colitis. *J Crohns Colitis*. 2020 May;14(4):465–79.
- 5 Calabrese E, Maaser C, Zorzi F, Kannengiesser K, Hanauer SB, Bruining DH, et al. Bowel Ultrasonography in the Management of Crohn's Disease. A Review with Recommendations of an International Panel of Experts. *Inflamm Bowel Dis*. 2016 May;22(5):1168–83.
- 6 de Voogd FA, Verstockt B, Maaser C, Gece KB. Point-of-care intestinal ultrasonography in inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol*. 2021 Apr;18(4):209–10.
- 7 Goodsall TM, Nguyen TM, Parker CE, Ma C, Andrews JM, Jairath V, et al. Systematic Review: Gastrointestinal Ultrasound Scoring Indices for Inflammatory Bowel Disease. *J Crohns Colitis*. 2021 Jan;15(1):125–42.
- 8 Astegiano M, Bresso F, Cammarota T, Sarno A, Robotti D, Demarchi B, et al. Abdominal pain and bowel dysfunction: diagnostic role of intestinal ultrasound. *Eur J Gastroenterol Hepatol*. 2001 Aug;13(8):927–31.
- 9 White LS, Campbell C, Lee A, Lord A, Graham RS. Intestinal ultrasound as first-line investigation in low risk gastrointestinal symptoms: a new model of care. *Intern Med J*. 2020 Nov;16:imj.15133.
- 10 Revzin MV, Moshiri M, Bokhari J, Pellerito JS, Menias C. Sonographic assessment of infectious diseases of the gastrointestinal tract: from scanning to diagnosis. *Abdom Radiol*. 2020 Feb;45(2):261–92.
- 11 Kucharzik T, Kannengiesser K, Petersen F. The use of ultrasound in inflammatory bowel disease. *Ann Gastroenterol*. 2017;30(2):135–44.
- 12 Dong J, Wang H, Zhao J, Zhu W, Zhang L, Gong J, et al. Ultrasound as a diagnostic tool in detecting active Crohn's disease: a meta-analysis of prospective studies. *Eur Radiol*. 2014 Jan;24(1):26–33.
- 13 Bollegala N, Griller N, Bannerman H, Habal M, Nguyen GC. Ultrasound vs Endoscopy, Surgery, or Pathology for the Diagnosis of Small Bowel Crohn's Disease and its Complications. *Inflamm Bowel Dis*. 2019 Jul;25(8):1313–38.
- 14 Ellrichmann M, Wietzke-Braun P, Dhar S, Nikolaus S, Arlt A, Bethge J, et al. Endoscopic ultrasound of the colon for the differentiation of Crohn's disease and ulcerative colitis in comparison with healthy controls. *Aliment Pharmacol Ther*. 2014 Apr;39(8):823–33.
- 15 Panés J, Bouzas R, Chaparro M, García-Sánchez V, Gisbert JP, Martínez de Guereñu B, et al. Systematic review: the use of ultrasonography, computed tomography and magnetic resonance imaging for the diagnosis, assessment of activity and abdominal complications of Crohn's disease. *Aliment Pharmacol Ther*. 2011 Jul;34(2):125–45.
- 16 Allocca M, Fiorino G, Bonifacio C, Furfaro F, Gilardi D, Argollo M, et al. Comparative Accuracy of Bowel Ultrasound Versus Magnetic Resonance Enterography in Combination With Colonoscopy in Assessing Crohn's Disease and Guiding Clinical Decision-making. *J Crohns Colitis*. 2018 Nov;12(11):1280–7.
- 17 Yuksel I, Kilincalp S, Coskun Y, Akinci H, Hamamci M, Alkan A. Diagnostic accuracy of intestinal ultrasound and magnetic resonance enterography for the detection of endoscopy-based disease activity in ileocolonic Crohn's disease. *Eur J Gastroenterol Hepatol*. 2019 Jul;31(7):809–16.
- 18 Taylor SA, Mallett S, Bhatnagar G, Baldwin-Cleland R, Bloom S, Gupta A, et al.; METRIC study investigators. Diagnostic accuracy of magnetic resonance enterography and small bowel ultrasound for the extent and activity of newly diagnosed and relapsed Crohn's disease (METRIC): a multicentre trial. *Lancet Gastroenterol Hepatol*. 2018 Aug;3(8):548–58.
- 19 Novak K, Maaser C, Wilkens R, Maconi G, Kucharzik T; International Bowel Ultrasound Group. Diagnostic accuracy of MRE and ultrasound for Crohn's disease. *Lancet Gastroenterol Hepatol*. 2019 Feb;4(2):95–6.
- 20 Sturm A, Maaser C, Calabrese E, Annesse V, Fiorino G, Kucharzik T, et al.; European Crohn's and Colitis Organisation [ECCO] and the European Society of Gastrointestinal and Abdominal Radiology [ESGAR]. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 2: IBD scores and general principles and technical aspects. *J Crohns Colitis*. 2019 Mar;13(3):273–84.
- 21 Rispo A, Imperatore N, Testa A, Nardone OM, Luglio G, Caporaso N, et al. Diagnostic Accuracy of Ultrasonography in the Detection of Postsurgical Recurrence in Crohn's Disease: A Systematic Review with Meta-analysis. *Inflamm Bowel Dis*. 2018 Apr;24(5):977–88.
- 22 Paredes JM, Ripollés T, Cortés X, Moreno N, Martínez MJ, Bustamante-Balén M, et al. Contrast-enhanced ultrasonography: usefulness in the assessment of postoperative recurrence of Crohn's disease. *J Crohns Colitis*. 2013 Apr;7(3):192–201.
- 23 Calabrese E, Petruzzello C, Onali S, Condino G, Zorzi F, Pallone F, et al. Severity of postoperative recurrence in Crohn's disease: correlation between endoscopic and sonographic findings. *Inflamm Bowel Dis*. 2009 Nov;15(11):1635–42.
- 24 Losco A, Viganò C, Conte D, Cesana BM, Basilisco G. Assessing the activity of perianal Crohn's disease: comparison of clinical indices and computer-assisted anal ultrasound. *Inflamm Bowel Dis*. 2009 May;15(5):742–9.
- 25 Maconi G, Tonolini M, Monteleone M, Bezzio C, Furfaro F, Villa C, et al. Transperineal perineal ultrasound versus magnetic resonance imaging in the assessment of perianal Crohn's disease. *Inflamm Bowel Dis*. 2013 Dec;19(13):2737–43.
- 26 Bezzio C, Bryant RV, Manes G, Maconi G, Saibeni S. New horizons in the imaging of perianal Crohn's disease: transperineal ultrasonography. *Expert Rev Gastroenterol Hepatol*. 2017 Jun;11(6):523–30.
- 27 Wright EK, Novak KL, Lu C, Panaccione R, Ghosh S, Wilson SR. Transperineal ultrasonography in perianal Crohn disease: A valuable imaging modality. *Can J Gastroenterol Hepatol*. 2015 Nov-Dec;29(8):445–7.
- 28 Kakkadasam Ramaswamy P, Vizhi N K, Yel-sangikar A, Krishnamurthy AN, Bhat V, Bhat N. Utility of bowel ultrasound in assessing disease activity in Crohn's disease. *Indian J Gastroenterol*. 2020 Oct;39(5):495–502.
- 29 Futagami Y, Haruma K, Hata J, Fujimura J, Tani H, Okamoto E, et al. Development and validation of an ultrasonographic activity index of Crohn's disease. *Eur J Gastroenterol Hepatol*. 1999 Sep;11(9):1007–12.
- 30 Novak KL, Kaplan GG, Panaccione R, Afshar EE, Tanyingoh D, Swain M, et al. A Simple Ultrasound Score for the Accurate Detection of Inflammatory Activity in Crohn's Disease. *Inflamm Bowel Dis*. 2017 Nov;23(11):2001–10.
- 31 Neye H, Voderholzer W, Rickes S, Weber J, Wermke W, Lochs H. Evaluation of criteria for the activity of Crohn's disease by power Doppler sonography. *Dig Dis*. 2004;22(1):67–72.
- 32 Allocca M, Craviotto V, Bonovas S, Furfaro F, Zilli A, Peyrin-Biroulet L, et al. Predictive value of bowel ultrasound in Crohn's disease: a 12-month prospective study. *Clin Gastroenterol Hepatol*. 2021 Apr;22:S1542–3565.

- 33 Novak KL, Nylund K, Maaser C, Petersen F, Kucharzik T, Lu C, et al. Expert Consensus on Optimal Acquisition and Development of the International Bowel Ultrasound Segmental Activity Score [IBUS-SAS]: A Reliability and Inter-rater Variability Study on Intestinal Ultrasonography in Crohn's Disease. *J Crohns Colitis*. 2021 Apr;15(4):609–16.
- 34 Paredes JM, Ripollés T, Cortés X, Reyes MD, López A, Martínez MJ, et al. Non-invasive diagnosis and grading of postsurgical endoscopic recurrence in Crohn's disease: usefulness of abdominal ultrasonography and leucocyte scintigraphy. *J Crohns Colitis*. 2010;4(5):537–45.
- 35 Maconi G, Carsana L, Fociani P, Sampietro GM, Ardizzone S, Cristaldi M, et al. Small bowel stenosis in Crohn's disease: clinical, biochemical and ultrasonographic evaluation of histological features. *Aliment Pharmacol Ther*. 2003 Oct;18(7):749–56.
- 36 Lenze F, Wessling J, Bremer J, Ullerich H, Spieker T, Weckesser M, et al. Detection and differentiation of inflammatory versus fibromatous Crohn's disease strictures: prospective comparison of 18F-FDG-PET/CT, MR-enteroclysis, and transabdominal ultrasound versus endoscopic/histologic evaluation. *Inflamm Bowel Dis*. 2012 Dec;18(12):2252–60.
- 37 Rispo A, Imperatore N, Testa A, Mainenti P, De Palma GD, Luglio G, et al. Bowel Damage in Crohn's Disease: Direct Comparison of Ultrasonography-based and Magnetic Resonance-based Lemann Index. *Inflamm Bowel Dis*. 2017 Jan;23(1):143–51.
- 38 Fraquelli M, Castiglione F, Calabrese E, Maconi G. Impact of intestinal ultrasound on the management of patients with inflammatory bowel disease: how to apply scientific evidence to clinical practice. *Dig Liver Dis*. 2020 Jan;52(1):9–18.
- 39 Allocca M, Fiorino G, Bonovas S, Furfaro F, Gilardi D, Argollo M, et al. Accuracy of Humanitas Ultrasound Criteria in Assessing Disease Activity and Severity in Ulcerative Colitis: A Prospective Study. *J Crohns Colitis*. 2018 Nov;12(12):1385–91.
- 40 Pascu M, Roznowski AB, Müller HP, Adler A, Wiedenmann B, Dignass AU. Clinical relevance of transabdominal ultrasonography and magnetic resonance imaging in patients with inflammatory bowel disease of the terminal ileum and large bowel. *Inflamm Bowel Dis*. 2004 Jul;10(4):373–82.
- 41 Parente F, Greco S, Molteni M, Cucino C, Maconi G, Sampietro GM, et al. Role of early ultrasound in detecting inflammatory intestinal disorders and identifying their anatomical location within the bowel. *Aliment Pharmacol Ther*. 2003 Nov;18(10):1009–16.
- 42 Sagami S, Kobayashi T, Aihara K, Umeda M, Morikubo H, Matsubayashi M, et al. Transperineal ultrasound predicts endoscopic and histological healing in ulcerative colitis. *Aliment Pharmacol Ther*. 2020 Jun;51(12):1373–83.
- 43 Civitelli F, Di Nardo G, Oliva S, Nuti F, Ferrari F, Dilillo A, et al. Ultrasonography of the colon in pediatric ulcerative colitis: a prospective, blind, comparative study with colonoscopy. *J Pediatr*. 2014 Jul;165(1):78–84.e2.
- 44 Hashimoto Y, Kume N, Sato K, Kanemura T, Haga A, Ishii Y, et al. P331 Development of a novel transabdominal ultrasound disease activity score in patients with ulcerative colitis (UCUS score). *J Crohns Colitis*. 2018;12(suppl. 1):S269.
- 45 Parente F, Molteni M, Marino B, Colli A, Ardizzone S, Greco S, et al. Are colonoscopy and bowel ultrasound useful for assessing response to short-term therapy and predicting disease outcome of moderate-to-severe forms of ulcerative colitis?: a prospective study. *Am J Gastroenterol*. 2010 May;105(5):1150–7.
- 46 Ramaswamy KP, Yelsangikar A, Nagarajan KV, Nagar A, Bhat N. P154 Development of a novel ultrasound based score for assessing disease activity in ulcerative colitis: preliminary results. *J Crohns Colitis*. 2019;13(suppl. 1):S166.
- 47 Ishikawa D, Ando T, Watanabe O, Ishiguro K, Maeda O, Miyake N, et al. Images of colonic real-time tissue sonoelastography correlate with those of colonoscopy and may predict response to therapy in patients with ulcerative colitis. *BMCGastroenterol*. 2011 Mar;11(1):29.
- 48 Kinoshita K, Katsurata T, Nishida M, Omotahara S, Onishi R, Mabe K, et al. Usefulness of transabdominal ultrasonography for assessing ulcerative colitis: a prospective, multicenter study. *J Gastroenterol*. 2019 Jun;54(6):521–9.
- 49 Maconi G, Bollani S, Bianchi Porro G. Ultrasonographic detection of intestinal complications in Crohn's disease. *Dig Dis Sci*. 1996 Aug;41(8):1643–8.
- 50 Kohn A, Cerro P, Milite G, De Angelis E, Prantera C. Prospective evaluation of transabdominal bowel sonography in the diagnosis of intestinal obstruction in Crohn's disease: comparison with plain abdominal film and small bowel enteroclysis. *Inflamm Bowel Dis*. 1999 Aug;5(3):153–7.
- 51 Gasche C, Moser G, Turetschek K, Schober E, Moeschl P, Oberhuber G. Transabdominal bowel sonography for the detection of intestinal complications in Crohn's disease. *Gut*. 1999 Jan;44(1):112–7.
- 52 Neye H, Ensberg D, Rauh P, Peitz U, Mönkemüller K, Treiber G, et al. Impact of high-resolution transabdominal ultrasound in the diagnosis of complications of Crohn's disease. *Scand J Gastroenterol*. 2010 Jun;45(6):690–5.
- 53 Parente F, Maconi G, Bollani S, Anderloni A, Sampietro G, Cristaldi M, et al. Bowel ultrasound in assessment of Crohn's disease and detection of related small bowel strictures: a prospective comparative study versus X-ray and intraoperative findings. *Gut*. 2002 Apr;50(4):490–5.
- 54 Kucharzik T, Maaser C. Intestinal ultrasound and management of small bowel Crohn's disease. *Therap Adv Gastroenterol*. 2018 May;11:1756284818771367.
- 55 Castiglione F, Mainenti PP, De Palma GD, Testa A, Bucci L, Pesce G, et al. Noninvasive diagnosis of small bowel Crohn's disease: direct comparison of bowel sonography and magnetic resonance enterography. *Inflamm Bowel Dis*. 2013 Apr;19(5):991–8.
- 56 Calabrese E, La Seta F, Buccellato A, Virdone R, Pallotta N, Corazziari E, et al. Crohn's disease: a comparative prospective study of transabdominal ultrasonography, small intestine contrast ultrasonography, and small bowel enema. *Inflamm Bowel Dis*. 2005 Feb;11(2):139–45.
- 57 Parente F, Greco S, Molteni M, Anderloni A, Sampietro GM, Danelli PG, et al. Oral contrast enhanced bowel ultrasonography in the assessment of small intestine Crohn's disease. A prospective comparison with conventional ultrasound, x ray studies, and ileocolonoscopy. *Gut*. 2004 Nov;53(11):1652–7.
- 58 Pallotta N, Vincoli G, Montesani C, Chirletti P, Pronio A, Caronna R, et al. Small intestine contrast ultrasonography (SICUS) for the detection of small bowel complications in crohn's disease: a prospective comparative study versus intraoperative findings. *Inflamm Bowel Dis*. 2012 Jan;18(1):74–84.
- 59 Calabrese E, Zorzi F, Onali S, Stasi E, Fiori R, Prencipe S, et al. Accuracy of small-intestine contrast ultrasonography, compared with computed tomography enteroclysis, in characterizing lesions in patients with Crohn's disease. *Clin Gastroenterol Hepatol*. 2013 Aug;11(8):950–5.
- 60 Ripollés T, Rausell N, Paredes JM, Grau E, Martínez MJ, Vizúete J. Effectiveness of contrast-enhanced ultrasound for characterization of intestinal inflammation in Crohn's disease: a comparison with surgical histopathology analysis. *J Crohns Colitis*. 2013 Mar;7(2):120–8.
- 61 Bettenworth D, Bokemeyer A, Baker M, Mao R, Parker CE, Nguyen T, et al.; Stenosis Therapy and Anti-Fibrotic Research (STAR) Consortium. Assessment of Crohn's disease-associated small bowel strictures and fibrosis on cross-sectional imaging: a systematic review. *Gut*. 2019 Jun;68(6):1115–26.
- 62 Ripollés T, Martínez-Pérez MJ, Paredes JM, Vizúete J, García-Martínez E, Jiménez-Restrepo DH. Contrast-enhanced ultrasound in the differentiation between phlegmon and abscess in Crohn's disease and other abdominal conditions. *Eur J Radiol*. 2013 Oct;82(10):e525–31.
- 63 Esteban JM, Aleixandre A, Hurtado MJ, Maldonado L, Mora FJ, Nogués E. Contrast-enhanced power Doppler ultrasound in the diagnosis and follow-up of inflammatory abdominal masses in Crohn's disease. *Eur J Gastroenterol Hepatol*. 2003 Mar;15(3):253–9.
- 64 Harbord M, Eliakim R, Bettenworth D, Karmiris K, Katsanos K, Kopylov U, et al.; European Crohn's and Colitis Organisation [ECCO]. Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 2: current Management. *J Crohns Colitis*. 2017 Jul;11(7):769–84.

- 65 Arienti V, Campieri M, Boriani L, Gionchetti P, Califano C, Giancane S, et al. Management of severe ulcerative colitis with the help of high resolution ultrasonography. *Am J Gastroenterol*. 1996 Oct;91(10):2163–9.
- 66 Smith R, Taylor K, Friedman A, Swaine A, Gibson D, Gibson P. Early assessment with gastrointestinal ultrasound in patients hospitalised for a flare of ulcerative colitis and predicting the need for salvage therapy: a pilot study. *Ultrasound Med Biol*. 2021;(5):S0301–5629–3.
- 67 Scarallo L, Maniscalco V, Paci M, Renzo S, Naldini S, Barp J, et al. Bowel Ultrasound Scan Predicts Corticosteroid Failure in Children With Acute Severe Colitis. *J Pediatr Gastroenterol Nutr*. 2020 Jul;71(1):46–51.
- 68 Ripollés T, Paredes JM, Martínez-Pérez MJ, Rimola J, Jauregui-Amezaga A, Bouzas R, et al. Ultrasonographic Changes at 12 Weeks of Anti-TNF Drugs Predict 1-year Sonographic Response and Clinical Outcome in Crohn's Disease: A Multicenter Study. *Inflamm Bowel Dis*. 2016 Oct;22(10):2465–73.
- 69 Maconi G, Ardizzone S, Parente F, Bianchi Porro G. Ultrasonography in the evaluation of extension, activity, and follow-up of ulcerative colitis. *Scand J Gastroenterol*. 1999 Nov;34(11):1103–7.
- 70 Kucharzik T, Wittig BM, Helwig U, Börner N, Rössler A, Rath S, et al.; TRUST study group. Use of Intestinal Ultrasound to Monitor Crohn's Disease Activity. *Clin Gastroenterol Hepatol*. 2017 Apr;15(4):535–542.e2.
- 71 Paredes JM, Ripollés T, Cortés X, Martínez MJ, Barrachina M, Gómez F, et al. Abdominal sonographic changes after antibody to tumor necrosis factor (anti-TNF) alpha therapy in Crohn's Disease. *Dig Dis Sci*. 2010 Feb;55(2):404–10.
- 72 Castiglione F, Imperatore N, Testa A, De Palma GD, Nardone OM, Pellegrini L, et al. One-year clinical outcomes with biologics in Crohn's disease: transmural healing compared with mucosal or no healing. *Aliment Pharmacol Ther*. 2019 Apr;49(8):1026–39.
- 73 Kucharzik T, Wilkens R, Maconi G, D'Agostino MA, Le Bars M, Nazar M, et al. DOP10 Intestinal ultrasound response and transmural healing after ustekinumab induction in Crohn's disease: Week 16 interim analysis of the STARDUST trial substudy. *J Crohns Colitis*. 2020;14 Suppl 1:S046–8.
- 74 Kucharzik T, Wilkens R, Maconi G, et al. LB12 Intestinal ultrasound response and transmural healing after 48 weeks of treatment with ustekinumab in Crohn's disease: STARDUST trial substudy. *United European Gastroenterol J*. 2020;8(10):1258–75.
- 75 Calabrese E, Rispo A, Zorzi F, et al. Ultrasonography tight control and monitoring in Crohn's disease during different biological therapies: a multicenter study. *Clin Gastroenterol Hepatol*. 2021;21:S1542–3565.
- 76 Zorzi F, Ghosh S, Chiamonte C, Lolli E, Ventura M, Onali S, et al. Response Assessed by Ultrasonography as Target of Biological Treatment for Crohn's Disease. *Clin Gastroenterol Hepatol*. 2020 Aug;18(9):2030–7.
- 77 Quaia E, Gennari AG, Cova MA. Early Predictors of the Long-term Response to Therapy in Patients With Crohn Disease Derived From a Time-Intensity Curve Analysis After Microbubble Contrast Agent Injection. *J Ultrasound Med*. 2019 Apr;38(4):947–58.
- 78 Goertz RS, Klett D, Wildner D, Atreya R, Neurath MF, Strobel D. Quantitative contrast-enhanced ultrasound for monitoring vedolizumab therapy in inflammatory bowel disease patients: a pilot study. *Acta Radiol*. 2018 Oct;59(10):1149–56.
- 79 Maaser C, Petersen F, Helwig U, Fischer I, Roessler A, Rath S, et al.; German IBD Study Group and the TRUST&UC study group; German IBD Study Group and TRUST&UC study group. Intestinal ultrasound for monitoring therapeutic response in patients with ulcerative colitis: results from the TRUST&UC study. *Gut*. 2020 Sep;69(9):1629–36.
- 80 Fraquelli M, Sarno A, Girelli C, Laudi C, Buscarini E, Villa C, et al. Reproducibility of bowel ultrasonography in the evaluation of Crohn's disease. *Dig Liver Dis*. 2008 Nov;40(11):860–6.
- 81 De Voogd F, Wilkens R, Gecse K, Allocca M, Novak K, Lu C, et al. A reliability study: strong inter-observer agreement of an expert panel for intestinal ultrasound in ulcerative colitis. *J Crohns Colitis*. 2021 Aug;15(8):1284–90.
- 82 Calabrese E. Bowel ultrasound for the assessment of Crohn's disease. *Gastroenterol Hepatol (N Y)*. 2011 Feb;7(2):107–9.
- 83 Bots S, Nylund K, Lowenberg M, Gecse K, D'Haens G. Intestinal ultrasound to assess disease activity in ulcerative colitis: development of a novel UC-ultrasound index. *J Crohns Colitis*. 2021;7(8):1264–71.
- 84 Omotehara S, Nishida M, Kinoshita K, et al. Validation of US evaluation of ulcerative colitis activity. *Ultrasound Med Biol*. 2019; 45(7):1537–44.