

# Lower Gastrointestinal Bleeding after Gynecological Surgery: An Atypical Endoscopic Diagnosis

Vítor Macedo Silva<sup>a, b, c</sup> Tiago Lima Capela<sup>a, b, c</sup> Pedro Boal Carvalho<sup>a, b, c</sup>  
Bruno Rosa<sup>a, b, c</sup> José Cotter<sup>a, b, c</sup>

<sup>a</sup>Gastroenterology Department, Hospital da Senhora da Oliveira, Guimaraes, Portugal; <sup>b</sup>Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal; <sup>c</sup>ICVS/3B's – PT Government Associate Laboratory, Guimaraes, Portugal

## Keywords

Lower gastrointestinal bleeding · Surgical trauma · Rectum

**Hemorragia Digestiva Baixa Após Cirurgia Ginecológica: Um Diagnóstico Endoscópico Atípico**

## Palavras Chave

Hemorragia digestiva baixa · Trauma cirúrgico · Recto


A 75-year-old woman with arterial hypertension and dyslipidemia was hospitalized in the Gynecology Department after transvaginal hysterectomy with anterior colporrhaphy due to grade IV hysterocele and cystocele. One week after the surgical procedure, the patient developed constipation, associated with abundant rectal bleeding and hypotension (90/55 mm Hg), without as-

sociated tachycardia. The patient reported no abdominal pain. There was no history of antiplatelet or anticoagulant agents' usage.


Analytically, a de novo microcytic anemia was detected, with hemoglobin levels of 8.4 g/dL (previously available value was 12.2 g/dL). Coagulation tests were normal.

In the setting of an acute lower gastrointestinal bleeding, an urgent colonoscopy after bowel preparation was conducted. Proctologic examination was normal. On digital rectal examination, a soft bulge was felt on the distal rectum, with bright blood being detected.

On colonoscopy, a voluminous bulge (approximately 60 × 55 mm) with a congestive appearance was seen in the distal rectum, with a 10-mm orifice being evident on its surface (Fig. 1). Inside this orifice, there was abundant coagulated hematic content, which could not be aspirated. This bulge was passable by the



**Fig. 1.** Endoscopic images showing a voluminous bulge (approximately 60 × 55 mm) with a congestive appearance seen in the distal rectum, with a 10-mm orifice being evident on its surface, compatible with a rectal hematoma.




**Fig. 2.** Abdominal and pelvic computerized tomography revealing an intramural collection containing hydro-aeric levels in the right rectal wall, with a maximum diameter of 7 cm, compressing the rectal lumen.

conventional colonoscope, without resistance. There were no other relevant endoscopic findings throughout the colon.

An abdominal and pelvic computerized tomography was performed, revealing an intramural collection containing hydro-aeric levels in the right rectal wall, with a maximum diameter of 7 cm, compressing the rectal lumen (Fig. 2). The lower gastrointestinal bleeding and endoscopic findings were thus interpreted in the context of a rectal mural hematoma, communicating with the rectal lumen. The patient underwent conservative treatment, with complete reabsorption of the hematoma 2 months after surgery, which was confirmed by cross-sectional imaging and colonoscopy (Fig. 3).

Small pelvic hematomas are a common finding after transvaginal hysterectomy, occurring in more than 25% of the performed surgeries [1]. Nevertheless, most of these are asymptomatic, being accidentally detected in scheduled follow-up ultrasounds. Furthermore, hematomas most frequently form in dependent areas, such as the pouch of Douglas, sub-vesical space, or ischium-rectal fossa [2]. Treatment consists of antibiotics if infected, with or without percutaneous or surgical drainage according to clinical evolution [1].

To our knowledge, this is the first case reporting a symptomatic rectal hematoma after gynecological surgery. With this clinical case, we aim to highlight an unusual etiology of lower gastrointestinal bleeding, associated with unique endoscopic findings, as well as to emphasize the importance of investigating medical and surgical history in patients with acute rectal bleeding.



**Fig. 3.** Colonoscopy (a) and computed tomography (b) images confirming complete rectal hematoma resolution.

### Statement of Ethics

Written informed consent was obtained from the patient for publication of the details of their medical case and any accompanying images. Ethical Committee approval was not required for this study, in accordance with local/national guidelines.

### Conflict of Interest Statement

The authors have no conflicts of interest to declare.

### Funding Sources

There was no funding for this work.

### References

- 1 Chill HH, Amsalem I, Karavani G, Amit S, Benshushan A, Shveiky D. Symptomatic pelvic hematoma following hysterectomy: risk factors, bacterial pathogens and clinical outcome. *BMC Womens Health*. 2020;20(1):272.
- 2 Alhubaishi L, Fazari A. Vault hematoma after vaginal hysterectomy: an invariable complication. *Open J Obstet Gynecol*. 2019;09(05): 578–81.

### Author Contributions

Vítor Macedo Silva, Tiago Lima Capela, and Bruno Rosa were involved in the endoscopic procedure and in manuscript drafting. Pedro Boal Carvalho and Bruno Rosa were involved in manuscript drafting and critical revision. José Cotter reviewed the manuscript and gave final approval. All authors approved the final version.

### Data Availability Statement

All data generated or analyzed during this study are included in this article. Further inquiries can be directed to the corresponding author.