CIÊNCIAS DA VIDA E DA SAÚDE LIFE AND HEALTH SCIENCES CIENCIAS DE LA VIDA Y LA SALUD

millenium

Millenium, 2(20)

PERFIL DE STRESS DOS ENFERMEIROS DE URGÊNCIA/EMERGÊNCIA EM CONTEXTO COVID-19: DETERMINANTES PROFISSIONAIS

STRESS PROFILE OF URGENCY/EMERGENCY NURSES IN COVID-19 CONTEXT: PROFESSIONAL DETERMINANTS

PERFIL DE ESTRÉS DE LAS ENFERMERAS DE URGENCIA/EMERGENCIA EN EL CONTEXTO COVID-19: DETERMINANTES PROFESIONALES

Nuno Duarte Pinto^{1,2} ^(D) https://orcid.org/0000-0003-3566-0538 Vitor Alves^{1,2} ^(D) https://orcid.org/0000-0003-0657-4745 Maria Augusta Veiga-Branco¹ ^(D) https://orcid.org/0000-0002-7963-2291

¹ Instituto Politécnico de Bragança, Bragança, Portugal
² Instituto Nacional de Emergência Médica, Moimenta da Beira, Portugal

Nuno Duarte Pinto - nunoduartepinto@gmail.com | Vitor Alves - vitor_alves940@hotmail.com | Maria Augusta Veiga-Branco - aubra@ipb.pt



Corresponding Author Nuno Duarte Pinto Rua do Canedo, N.28, Oliveira de Barreiros 3500-892 – Viseu – Portugal nunoduartepinto@gmail.com RECEIVED: 12th July, 2022 REVIEWED: 15th November, 2022 ACCEPTED: 07th December, 2022 PUBLISHED: 31th January, 2023 Pinto, N. D., Alves, V., & Veiga-Branco, M. A. (2023). Stress profile of urgency/emergency nurses in covid-19 context: professional determinants. *Millenium - Journal of Education, Technologies, and Health, 2*(20), 1-10, e27652. **DOI**: https://doi.org/10.29352/mill0220.27652

RESUMO

Introdução: As evidências revelam que os enfermeiros de urgência/emergência desenvolveram problemas de saúde relacionados com o *stress* em contexto de pandemia por COVID-19.

Objetivos: Avaliar o perfil de *stress* dos enfermeiros de urgência/emergência em contexto COVID-19 em função das variáveis profissionais.

Métodos: Estudo transversal, quantitativo, descritivo-correlacional, através da aplicação da Escala de Perceção de *Stress* (EPS), adaptada por Pais Ribeiro & Marques (2009), a uma amostra de 355 enfermeiros portugueses, em contexto de urgência/emergência.

Resultados: Maioritariamente é uma amostra feminina (51,1%), com idades entre 36-50 anos (51,3%), licenciatura (38,9%), exercem funções no setor público (85,9%), no Norte (35,8%), com tempo de experiência profissional entre 11-20 anos (40,8%) e com a categoria profissional de "enfermeiro" (59,4%).

Conclusão: As variáveis de caracterização profissional, revelaram que o título profissional, o serviço/unidade onde exercem funções e a localização geográfica, apresentam efeito significativo no perfil de *Stress* percecionado (p < 0,005). Neste âmbito, os enfermeiros especialistas (p = 0,005) a exercerem funções nos SUP e BO (p = 0,002) da Região de Lisboa e Vale do Tejo (p = 0,001), foram aqueles onde a perceção de *Stress* é mais alta. Por outro lado, os enfermeiros generalistas (p = 0,005), que exercem funções nos INEM (p = 0,002), e que exercem na Região Autónoma da Madeira (p = 0,001) demonstraram um perfil de *Stress* mais baixo.

Palavras-chave: stress ocupacional; enfermeiros; pandemia COVID-19; urgência; emergência

ABSTRACT

Introduction: Evidence shows that emergency nurses have developed *stress*-related health problems in the context of the COVID-19 pandemic.

Objectives: Evaluate the *stress* profile of urgency/emergency nurses in a COVID-19 context according to professional variables. **Methods:** Cross-sectional, quantitative descriptive-correlational study, through the application of the *Stress* Perception Scale (SPS) adapted by Pais Bibeiro & Margues (2000) to a sample of 255 Perturbation purses in an emergency/urgency sector.

(SPS), adapted by Pais Ribeiro & Marques (2009), to a sample of 355 Portuguese nurses in an emergency/ urgency context. **Results:** The sample was mainly composed of women (51.1%), aged between 36-50 years (51.3%), with a university degree (38.9%), working in the public sector (85.9%), in the North (35.8%), between 11-20 years of professional experience (40.8%) and with the professional category of "nurse" (59.4%).

Conclusion: As variáveis de caracterização profissional, revelaram que o título profissional, o serviço/unidade onde exercem funções e a localização geográfica, apresentam efeito significativo no perfil de *Stress* perceived (p < 0.005). In this context, the specialist nurses (p = 0.005) working in the SUP and OR (p = 0.002) of the Lisbon and Tagus Valley Region (p = 0.001) were those with the highest perception of *Stress*. On the other hand, generalist nurses (p = 0.005), who exercise functions at INEM (p = 0.002), and who exercise in the Autonomous Region of Madeira (p = 0.001) showed a lower *Stress* profile.

Keywords: occupational stress; nurses; pandemic COVID-19; urgency; emergency

RESUMEN

Introducción: Los datos demuestran que las enfermeras de urgencias han desarrollado problemas de salud relacionados con el *estrés* en el contexto de la pandemia de COVID-19.

Objetivos: Evaluar el perfil de estrés de las enfermeras de urgencias/emergencias en un contexto COVID-19 según variables.

Métodos: Estudio transversal, cuantitativo, descriptivo y correlacional, a través de la aplicación de la Escala de Percepción de Estrés (EPS), adaptada por Pais Ribeiro & Marques (2009), a una muestra de 355 enfermeras portuguesas en un contexto de emergencia/urgencia.

Resultados: La muestra se compone principalmente de mujeres (51,1%), con edades comprendidas entre los 36 y los 50 años (51,3%), con titulación universitaria (38,9%), que trabajan en el sector público (85,9%), en el Norte (35,8%), entre 11 y 20 años de experiencia profesional (40,8%) y con la categoría profesional de "enfermera" (59,4%).

Conclusión: Las variables de caracterización profesional, revelaron que el título profesional, el servicio/unidad donde se ejercen las funciones y la localización geográfica, presentan un efecto significativo en el perfil de *Estrés* percibido (p < 0,005). En este contexto, los enfermeros especialistas (p = 0,005) que trabajan en el SUP y en el quirófano (p = 0,002) de la Región de Lisboa y del Valle del Tajo (p = 0,001) fueron los que tuvieron una mayor percepción de *Estrés*. Por otro lado, los enfermeros generalistas (p = 0,005), que ejercen funciones en el INEM (p = 0,002), y que ejercen en la Región Autónoma de Madeira (p = 0,001) mostraron un perfil de *Estrés* menor.

Palabras clave: estrés profesional; enfermeras; pandemia COVID-19; urgencia; emergencia



INTRODUCTION

Coronavirus, known as COVID-19, was identified in December of 2019 in China, in the city of Wuhan and on the 11th of March of 2020 the World Health Organization (WHO) assumed that this outbreak had rapidly turned into a global pandemic (Hu et al., 2020; He et al., 2020). The development of the dissemination originated that Portugal was no exception and the competent entities were obliged to implement restrictive and blocking measures, in an attempt to flatten the contagion curve, similarly to the rest of the world.

The interest in studying the stress profile of Portuguese urgency/emergency Nurses in the context of COVID-19, considering his socio-demographic and professional determinants, arises from the fact of having knowledge of the results of the study "Mental Health In Time of Pandemic" (MH-COVID-19)" (Almeida, et al., 2020), coordinated by the Health Promotion and Non-Transmittable Diseases Department of the Doutor Ricardo Jorge National Health Institute, in collaboration with the Environmental Health Institute of the Medicine Faculty of Lisbon University and with the Portuguese Society of Psychiatry and Mental Health. This study analysed the impact of the COVID-19 pandemic in mental health and wellbeing of the population in general and, of the health professionals in particularly, considering dimensions like anxiety, depression, post-traumatic stress, burnout and resilience, among others.

The pandemic has exacerbated stressors in healthcare systems or services, where burnout among nurses in response to workplace stress is already an epidemic (Duarte t al., 2020).

According recent studies, some health professionals, including nurses who work in the urgency/emergency context, developed psychological suffering, fatigue and exhaustion, while facing the COVID-19 pandemic (Albott et al., 2020; Walton et al., 2020. These health professionals "often face work overload, nurse/patient ratio disparity, resources limitations and imbalance between work and family life, affecting their mental health" (Nunes, 2019, p. 30).

In order to know which socio-demographic and professional variables influence the stress perception level of urgency/emergency nurses in COVID-19 context, was formulated the question "The socio-demographic and professional variables influence the stress perception level of urgency/emergency nurses in COVID-19 context", in an attempt to find any statistically significant difference between the variables in question.

1. THEORETICAL FRAMEWORK

At the end of 2019, WHO received alerts of a number of cases of pneumonia in the city of Wuhan, China. It was the starting point of an outbreak generated by a new coronavirus (SARS-CoV2), originator of the coronavirus disease (COVID-19) that, since then, has spread with increasing numbers of cases in other regions of the world. On the 30th of January of 2020, WHO declared a Public Health Emergency of international importance and, on the 11th of March of 2020, the disease became a pandemic (Harapan et al., 2020).

The public health emergencies, although evidencing the fundamental role of the National Health System in the containment of COVID-19, expose the structural deficiencies of the system, particularly, the lack and/or the uneven distribution of the health professionals and of the infrastructures of the medium and high complexity care. Thus, the interaction between the work environment and the human factors seem to have triggered physical and emotional responses in the health professionals, in the specific case of urgency/emergency nurses, which caused a decrease in their mental health and, consequently, of their quality of life.

In the urgency/emergency environment, it is common to find different complementary stressors. The natural stress required to the nurses to give care to the person in critical situation and their relatives, with the complexity that permeates the possibilities of intervention, may turn these health professionals more vulnerable to stress. Nurses deal daily with pain and suffering and are subjected to intense rhythms and long work hours, complex human relationship, lack of materials and a reduced number of professionals (Teixeira et al., 2021).

The COVID-19 pandemic caught the world's attention, specially the consequences of the high risk of stress. The majority of the nurses experiences moderate to elevated levels of stress because these professionals are particularly at risk of becoming infected with COVID-19 and by the healthcare system overload. The nurses that are on the front line, particularly the ones that work in urgency/emergency context, experience an overload of work and psychological stress (Oktovin & Peni, 2021). The complaints, like physical and psychological fatigue are frequently felt due to high number of people infected with COVID-19.

Temsah et al., (2020) refer that pandemic lead to a phenomena without precedents of psychological stress in nurses. The pandemic infectious diseases, like COVID-19, impose a significant level of anxiety and stress in health professionals that take care of infected people, whose main concern is the risk of transmitting the infection to their relatives or develop the disease themselves.

The authors above mentioned refer that usually, nurses are at risk of exposure to highly infectious pathogenic factors in care giving, not only for the exposure to the environment of the person infected or in the contact with biological samples, but also in the lack of Individual Protection Equipment (IPE), which makes them experience high levels of psychological stress. These also

68

need to deal with the unpredictability of their work schedule and higher workload, which oblige them to a bigger adjustment of their private and social life.

The reports of the social media document extreme exhaustion, physical discomfort of long work hours with facial masks and other IPE, fear of contagion and anguish in nurses. This combination of physical and emotional wear in a nursing work force already stressed became a mark of the COVID-19 pandemic (Arnetz, et al., 2020).

The consequences of the stress perception levels in the health of the nurses evaluated during the pandemic, must be seen through a perspective of health and occupational safety. The stress and the burnout were internationally recognized as works risks for the nurses (Pérez-Fuentes et al., 2018) before the pandemic.

Although the research suggests that occupational and personality factors play a role in stress, in 2019, WHO declared stress as an occupational phenomena, in addition of being a medical condition. The beginning of the COVID-19 pandemic increased in the middle of a nursing staff already tense, putting their mental health and well-being at risk (Lai et al., 2020).

2. METHODS

2.1 Sample

Quantitative, descriptive correlational study with a cross-sectional approach. 355 nurses who work in urgency/emergency context at national level participated. The sample was of the non-probabilistic, accidental or convenience type.

Data collection occurred between the 1st of February of 2021 and 28th of February of 2021, through the application of a questionnaire of socio-demographic and professional data (ad hoc), with validation of informed consent for the present study by the participants and the Stress Perception Scale (SPS), adapted for the Portuguese population by Pais Ribeiro & Marques (2009), with application authorization date of 10/20/2000, available via Google Forms, to all nurses nationwide.

Ethical principles were fulfilled. Study approved by the Ethics Commission of Polytechnic Institute of Bragança on the 12/11/2020 (Resolution no. 28/2020).

2.2 Data collection instrument

The data collection instrument includes the following measuring instruments:

- Questionnaire of socio-demographic and professional data (ad hoc);

- Stress Perception Scale (SPS), adapted for the Portuguese population by Pais Ribeiro & Marques (2009).

The SPS is, according the authors, a global measuring unit of Stress which proposes to evaluate the grade in which an individual appreciates his life situations like Stress (Cohen, Kamarck & Mermelstein, 1983 in Pais Ribeiro & Margues, 2009). The SPS scale, validated for the Portuguese population is a directional scale with 13 items that operationalizes the Stress Perception variable and that, from the way the sample responds, whose global score results from the sum of the values assigned to each one.

A higher score corresponds to greater perceived Stress in the last month. Being the maximum score of 52 points. There are no cut-off points. The items 4, 5, 6, 7, 9, 10 and 12 are inverted items (Pais Ribeiro & Margues, 2009).

2.3 Statistical analyses

For data analysis, descriptive statistics and analytical or inferential statistics were used. The descriptive analyses enabled to determine the absolute and relative frequencies, measures of central tendency, namely the medium and the dispersion measures like range of variation, the variation coefficient and the standard deviation.

For the inferential analyses was carried out in the first place, the normality test. By the Kolmogorov-Smirnov-Lilliefors (Chart 1), which was applied because, in this sample, with a dimension greater than 30 observations, it allowed to verify that the distribution of data referring to the dependent variable (Stress Perception) is framed within normality (p<0,05). Given this, and through the analysis of figure 1, the existence of a normal or close to normal distribution was assumed.

Chart 1-Kolmogorov-Smirnov-Lilliefors Normality Tes	st
---	----

			Kolmogorov-Sn	nirnov-Lillieforsª
			Statistics	р
Stress Perce	otion Scale		0,089	0,000***
*p<0,05	**p<0,01	***p<0,00		

p<0,05

Pinto, N. D., Alves, V., & Veiga-Branco, M. A. (2023). Stress profile of urgency/emergency nurses in covid-19 context: professional determinants. *Millenium - Journal of Education, Technologies, and Health, 2*(20), 1-10, e27652. DOI: https://doi.org/10.29352/mill0220.27652



Average = 41,36 Deviation of the Dev. = 4,207 N = 355

Figure 1 - Histogram of the Stress Perception Scale, with normality curve

In the inferential statistics, non parametric, morment tests were used: U Mann-Whitney test and Kruskall-Wallis test. The investigation questions were tested with a 95% of probability, from where results a level of significance of 5% (α =0,05). The decision criteria for the tests are based in the study of probabilities, confirming the investigation question if the probability is less than

0,05 and rejecting it if greater than this value. The following significance levels were used:

- $p \ge 0.05 not significant$
- p < 0.05 significant
- p < 0.01 quite significant
- p < 0.001 highly significant

All statistical treatment was processed using the SPSS program (Statistical Package for the Social Sciences) version 26.0 for Windows.

3. RESULTS

The sample was mostly female (n=181; 51%), predominantly aged between 36-50 years old (51,3%). Regarding the marital status, (48,7%) are married and in the academic qualifications prevailed the graduates with (38,9%).

	Total	
Variables	No. (355)	% (100.0)
Age	129	36.3
<35 years old	182	51.3
36-50 years old	43	12.1
51-65 years old	1	0.3
66-80 years old	_	-)-
Gender		
Feminine	181	51,0
Masculine	174	49,0
Marital status		
Single	100	28,2
Married	173	48,7
Non-marital partnership	59	16,6
Divorced	19	5,4
Widow	4	1,1
Academic qualifications		
Bachelor's degree	7	2,0
Licensed degree	138	38,9
Postgraduate studies	113	31,8
Master's degree	90	25,4
Doctorate	7	2,0

Chart 2 -	Socio-demograph	nic chara	cterization	of the	sample
01101101	Socio acinograpi	ne chara	CCCTILCCTOTT	or the	Janpie

participants with the professional category of "nurse" prevail (59,4%).

The majority of the elements of the sample works in the public sector (85,9%), in the north (35,8%), with professional experience time between 11-20 years (40,8%), ECPF-IP* employment relationship (58,6%), rotating work schedule (86,8%), <35 weekly hours (75,2%) and on a full-time job basis (98,9%). As for the service/unit where they work, non-assigned services dominate (22,3%) and

	T	otal
Variables	No.	%
	(355)	(100.0)
Type of management		
Public	305	85,9
Private	19	5,4
P-P Partnership	31	8,7
Geographical location		
North	127	35,8
Centre	114	32,1
Lisbon and Tagus Valley	85	23.9
Alenteio	10	2.8
Algarve	12	3.4
Azores	4	1 1
Madeira	3	0.8
Professional time	5	0,0
0-10 years	113	31.8
11-20 years	145	40.8
21-30 years	77	21 7
21-30 years	20	5.6
SI-40 years	20	5,0
	120	26.6
	130	1 1
	4	1,1
	208	20,0
CCFF-RRT*	7	2,0
Other*	Ь	1,7
Work schedule	40	12.0
Fixed	46	13,0
Rotating	308	86,8
Both	1	0,3
work regime	254	00.0
Full time	351	98,9
Part time	4	1,1
Weekly hours		
<35 hours	267	75,2
36-50 hours	74	20,8
>50 hours	14	3,9
Service/Unit		
OR*	14	3,9
NIME*	66	18,6
BES*	44	12,4
MS-US*	76	21,4
MES*	50	14,1
MICU*	26	7,3
Other	79	22,3
Professional title		
Nurse	211	59,4
Specialist nurse	144	40.6

Chart 3 - Professional Characterization of the sample

*ECPF-IP: Employment contract of public functions-Undetermined period |*IEC: Individual employment contract |*CLA: Collective Labour agreement | *ECPF-RRT: Employment contract of public functions-right resolutive term | *OR- Operating room | *NIME- National institute of medical emergency |*BES- Basic Emergency service | *MS-US: Medical-surgical urgency service | *MES: Multipurpose emergency service | *MICU: Multipurpose intensive care unit

MA



By analysing table 4, and with regard to the value of the responses given to the Stress Perception Scale (SPS), it is noted that in all 13 items the "Sometimes" answers dominate, with a score of 3. "Never", were the responses less referred in the first 12 items. As for the 13th item, the response that obtained the lowest frequency of responses was "Many Times".

			Responses		
Items	Never (0)	Almost Never (1)	Sometimes (2)	Frequently (3	Many Times (4)
Item 1 - In the last month, how often have you been upset about something that happened unexpectedly?	7	48	160	78	62
Item 2 - In the past month, how often have you felt that you were unable to control the things that are important in your life?	30	77	125	73	50
Item 3 - In the past month, how often have you felt nervous or stressed?	5	48	123	98	81
Item 4- In the last month, how often have you successfully tackled boring and boring things?	7	25	185	99	39
Item 5 In the past month, how often have you felt that you were effectively coping with major changes that were taking place in your life?	21	68	167	69	30
Item 6 - In the last month, how often have you felt confident in your ability to deal with your personal problems?	5	45	155	101	49
Item 7 - In the last month, how often did you feel that things were going your way?	13	97	166	65	14
Item 8 In the last month, how often have you noticed that you couldn't do all the things you had to do?	4	55	146	103	47
Item 9 - In the last month, how often have you felt able to control your irritations?	4	52	179	90	30
Item 10 - In the last month, how often did you feel that things were going for the best for you?	10	96	181	54	14
Item 11 In the last month, how often have you been irritated by things that happened that were beyond your control?	9	59	148	101	38
Item 12 - In the last month, how often have you been able to control your time?	6	91	160	72	26
Item 13 In the last month, how often did you feel that difficulties were piling up to the point where you were unable to overcome them?	39	119	138	43	16

Chart 4 - Stress perception (SP)

With the objective to know the influence of the type of management in the stress perception level, it was applied the Kruskal-Wallis test, whose results indicate a higher level of stress by the nurses that work in the private sector, being the ones that work in a public-private partnership institution those who manifest less perception of the stress, however, with the absence of different significant statistics (p>0,05).

Regarding the geographical location versus the stress perception in COVID-19 context, a Kruskal-Wallis test took place, from which a higher level of stress is highlighted by the nurses that work in Lisbon and Tagus Valley. The nurses that work in Madeira Autonomous Region are the ones that manifest less stress perception, with statistical differences quite significant (p=0,001).

As for the influence of professional practice time, a Kruskal-Wallis test was applied, whose data indicate a higher level of stress perception on the part of nurses with longer professional practice, followed by those with between 21-30 years and those with 11-20 years of service. Those with less years of service are those who manifest less perception of stress, however, with no statistically significant differences (p>0,05).

Finally, and with regard to the employment relationship, a Krustal-Wallis test was carried out, which highlights a higher level of stress perception on the part of nurses with a CIT*. Finally, and with regard to the employment relationship, a Krustal-Wallis test was carried out, which highlights a higher level of stress perception on the part of nurses with a CIT*. Finally, and with regard to the employment relationship, a Krustal-Wallis test was carried out, which highlights a higher level of stress perception on the part of nurses with a CIT*. Finally, and with regard to the employment relationship, a Krustal-Wallis test was carried out, which highlights a higher level of stress perception on the part of nurses with a CIT*. Finally, and with regard to the employment relationship, a Krustal-Wallis test was carried out, which highlights a higher level of stress perception on the part of nurses with a IEC; followed for those who possess ECPF-RRT* and a CLA*. Those who possess other type of employmet relationship manifest less perception of stress. However, also here with the absence of statistically significant differences (p>0,05).

Variables	Level of stress perception Average ordering
Type of management	
Public	180,22
Private	184,95
P-P Partnership	151,89
(p)	0,324
Geographical location	
North	155,69
Centre	182,20
Lisbon and Tagus Valley	215,12
Alentejo	186,35
Algarve	121,29
Azores	173,63
Madeira	116,17
(p)	0,001**
Professional time	
0-10 years	173,35
11-20 years	176,41
21-30 years	180,31
31-40 years	206,95
(p)	0,591
Professional relationship	
IEC*	195,31
CLA*	182,88
ECPF-IP*	168,03
ECPF-RRT*	186,14
Other	136,00
(p)	0,149
n<0.05 **n<0.01 ***n<0.001	

|--|

4. DISCUSSION

The socio-demographic and professional profile of a sample of 335 Portuguese nurses performing urgency/emergency duties indicates a sample mainly female (51,0%), with an average age of 39,25±8,8 years old, with the predominance of the age group of 36-50 years (51,3%). Most nurses are married (48,7%), licensed (38,9%), having an average professional time of 15,29±8,92 years and an average weekly working time of 39,39±6,08 hours. Most participants work in the public sector (85,9%) in the north (35,8%), following the ones who work in the centre area (32,1%) and in Lisbon and Tagus Valley (23,9%). Those with professional experience time between 11-20 years stand out, who have ECPF-IP* with 58,6%, practicing rotating hours, with \leq 35 hours/week. As for the service/unit where they work, in addition to non-assigned services (22,3%), those working at MS-US* and NIME* (21,4%; 18,6%, respectively) and with the professional category of "nurse" stand out (59,4%). These data, for the most part, corroborate the socio-demographic and professional profile of a sample of 453 nurses who participated in the study by Cui et al. (2021), where there was a prevalence of women (96,47%), but with a slightly lower mean age (M=33,15±8,38 years) and mean length of professional experience (M=11,33±9,25 years) compared to the nurses in the present study. In that study, the prevalence of licensed nurses (38.7%) and with an average weekly working time of 37.29±6.01 hours was also recorded, results corroborated in the present study.

Considering the objective of the current study, to verify if the socio-demographic (age, gender, marital status and academic qualifications) and professional (type of management, geographical location, professional time, professional relationship, schedule, work regime, weekly hours, service and professional title) variables influence the level of stress perception of the urgency/emergency nurses in COVID-19 context, it was stated that younger nurses (\leq 35 years old), male, divorced and the nurses with master's degree would be mainly subjected to stress. On the other hand, the geographical location where these work interfered on the stress perception, registering higher level of stress perception, with quite significant statistical differences (p=0,001). No statistically significant differences were recorded with regard to the type of management of the service location, length of professional practice and employment relationship. However, the average ranking values indicate a higher level of stress on the part of nurses working in the private sector, with longer professional experience, followed by those with between 21-30 years and by those with between 11-20 years of service and by nurses with an IEC*; followed by those with a ECPF-RRT* and a CLA*. In the studies found, nurses with less time in professional practice were those who manifested a higher level of perception of stress



(Jijun et al., 2020; Fernandez et al., 2020), which was not corroborated in the sample of the present study. These studies indicate that younger nurses are the most predisposed to stress, which is justified by the fact that younger individuals with less professional experience still do not have enough time to formulate effective coping strategies to deal with occupational stress and, therefore, they are more vulnerable.

According Urzal et al. (2021), the percentage of Portuguese nurses during the COVID-19 pandemic (20,4%) with symptoms of anxiety, stress, depression and post-traumatic stress disorder comes close to studies carried out in China in February (27,39%) and June (20,87%) of 2020 (Huang, et al., 2020; Zhang & Ma, Z.F., 2020). However, the use of different evaluation instruments in the several studies complicates a comparative analyses with the Portuguese reality in terms of mental health of the nurses during the pandemic. Pandilha and Silva (2020) state that the pandemic was at the origin of changes in the professional practice and in the personal life of nurses, with some of them even having to change residence during the pandemic, taking them away from their family life, which is corroborated in the present study, where the married or cohabiting nurses were more likely to develop higher levels of stress. The most evident psychological manifestations in the literature on nurses during the pandemic were higher levels of stress, exhaustion and depressive mood, strong psychological effects and difficulty in coping with the situation, high levels of anxiety, fear and psychological suffering (Fernandez et al., 2020; Zerbini et al., 2020; Aksoy & Koçak, 2020).

The COVID-19 pandemic has drawn the world's attention, especially with regard to the risk of stress. Most nurses must have experienced moderate to high levels of stress, as these professionals are particularly at risk of being infected with COVID-19 and by the burden on the healthcare system.

CONCLUSION

Nurses who work in an urgency/emergency context are the main health professionals in an epidemic and have the most contact with patients. These faced many challenges during the COVID-19 epidemic, being directly at risk, triggering higher levels of work stress (Zakeri et al., 021).

It was in this context, and taking into account personal experience, that the present study was developed, with a sample of 355 Portuguese nurses working in urgency/emergency, with an average age of 36,25±8,89 years. The evidence found allowed us to answer the research question, having concluded that the only professional variable with statistically significant interference in terms of the perception of stress of urgency/emergency nurses in the context of COVID was the geographical location where they work, being higher the level of stress of the nurses working in Lisbon and the Tagus Valley. It was also concluded that there was a higher level of stress on the nurses working in the private sector, with longer professional experience, followed by those with between 21-30 years and by those with between 11-20 years of service and on the nurses with an IEC*; followed by those who have a ECPF-RRT* and a CLA*, although without statistically significant relevance. Although the socio-demographic variables did not reveal statistically significant differences, it was found that younger nurses (<35 years old), male, divorced and nurses with a master's degree, were the ones who presented a greater perception of stress during the COVID-19 pandemic. A systematic review with a meta-analysis focusing on the prevalence of depression, anxiety and stress during COVID-19 concluded that almost a third of nurses had depression, anxiety and stress during the COVID-19 pandemic. Anxiety and depression were more reported among female nurses, while stress was more prevalent in male and younger nurses (Nadeem et al., 2021). The psychological effects of the pandemic situation on nurses must be urgently evaluated, understood and controlled. Understanding and assessing their effects are essential for promoting and protecting nurses' well-being and emotional resilience, as they can directly affect the quality of healthcare provided. Despite the growing scientific evidence about health professionals affected by high levels of stress that increased significantly during the pandemic, there was a shortage in the literature on the topic addressed in urgency/emergency nurses, which resulted in the main limitation of this study. Therefore, it is suggested that this study be replicated in order to understand how these professionals adapted to work-related stress during a pandemic, and it is also interesting to investigate their cognitive response after a pandemic crisis, in order to develop preventive measures to mitigate higher levels of stress and help in coping.

REFERENCES

- Aksoy, Y. E., & Koçak, V. (2020). Psychological effects of nurses and midwives due to COVID-19 outbreak: The case of Turkey. *Archives of Psychiatric Nursing*, 34(5), 427–433. https://doi.org/10.1016/j.apnu.2020.07.011
- Albott, C. S., Wozniak, J. R., McGlinch, B. P., Wall, M. H., Gold, B. S., & Vinogradov, S. (2020). Battle Buddies: Rapid Deployment of a Psychological Resilience Intervention for Health Care Workers During the Coronavirus Disease 2019 Pandemic. *Anesthesia and Analgesia*, 10.1213/ANE.000000000004912. https://doi.org/10.1213/ANE.00000000004912

- Almeida, T. C. de, Heitor, M. J., Santos, O., Costa, A., Virgolino, A., Rasga, C., Martiniano, H., & Vicente, A. (2020). Saúde mental em tempos de pandemia - SM-COVID-19: Relatório final (pp. 1–222) [Report]. Instituto Nacional de Saúde Doutor Ricardo Jorge, IP. https://repositorio.insa.pt/handle/10400.18/7245
- Arnetz, J. E., Goetz, C. M., Arnetz, B. B., & Arble, E. (2020). Nurse Reports of Stressful Situations during the COVID-19 Pandemic: Qualitative Analysis of Survey Responses. *International Journal of Environmental Research and Public Health*, *17*(21), 8126. https://doi.org/10.3390/ijerph17218126
- Duarte, I., Teixeira, A., Castro, L., Marina, S., Ribeiro, C., Jácome, C., Martins, V., Ribeiro-Vaz, I., Pinheiro, H. C., Silva, A. R., Ricou, M., Sousa, B., Alves, C., Oliveira, A., Silva, P., Nunes, R., & Serrão, C. (2020). Burnout among Portuguese healthcare workers during the COVID-19 pandemic. *BMC Public Health*, 20(1), 1885. https://doi.org/10.1186/s12889-020-09980-z
- Fernandez, R., Lord, H., Halcomb, E., Moxham, L., Middleton, R., Alananzeh, I., & Ellwood, L. (2020). Implications for COVID-19: A systematic review of nurses' experiences of working in acute care hospital settings during a respiratory pandemic. International Journal of Nursing Studies, 111, 103637. https://doi.org/10.1016/j.ijnurstu.2020.103637
- Harapan, H., Itoh, N., Yufika, A., Winardi, W., Keam, S., Te, H., Megawati, D., Hayati, Z., Wagner, A. L., & Mudatsir, M. (2020). Coronavirus disease 2019 (COVID-19): A literature review. *Journal of Infection and Public Health*, *13*(5), 667–673. https://doi.org/10.1016/j.jiph.2020.03.019
- He, F., Deng, Y., & Li, W. (2020). Coronavirus disease 2019: What we know? *Journal of Medical Virology*, *92*(7), 719–725. https://doi.org/10.1002/jmv.25766
- Hu, D., Kong, Y., Li, W., Han, Q., Zhang, X., Zhu, L. X., Wan, S. W., Liu, Z., Shen, Q., Yang, J., He, H.-G., & Zhu, J. (2020). Frontline nurses' burnout, anxiety, depression, and fear statuses and their associated factors during the COVID-19 outbreak in Wuhan, China: A large-scale cross-sectional study. *EClinicalMedicine*, 24, 100424. https://doi.org/10.1016/j.eclinm.2020.100424
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, *395*(10223), 497–506. https://doi.org/10.1016/S0140-6736(20)30183-5
- Jijun, W.; Xian, S.; Fei, C.; Yuanjie, D.; Dechun, C.; Xingcao, J., & Xiaoling, G. (2020). Survey of sleep quality and its influencing factors in clinical front-line nurses to combat the new coronavirus pneumonia epidemic. *Nursing Res.* 1–5. http://kns.cnki.net/kcms/detail/14.1272.r.20200214.1136.004.html.
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H., Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., & Hu, S. (2020). Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. JAMA Network Open, 3(3), e203976. https://doi.org/10.1001/jamanetworkopen.2020.3976
- Nadeem, F., Sadiq, A., Raziq, A., Iqbal, Q., Haider, S., Saleem, F., & Bashaar, M. (2021). Depression, Anxiety, and Stress Among Nurses During the COVID-19 Wave III: Results of a Cross-Sectional Assessment. *Journal of Multidisciplinary Healthcare*, 14, 3093–3101. https://doi.org/10.2147/JMDH.S338104
- Nunes, T. M. M. (2019). *Determinantes de burnout em enfermeiros do pré-hospitalar* [MasterThesis, Instituto Politécnico de Viseu. Escola Superior de Saúde de Viseu]. https://repositorio.ipv.pt/handle/10400.19/5547
- Oktovin, Basit, M., & Peni, M. R. (2021). Psychological Stress of Nurses During the Covid-19 Pandemic. *KnE Life Sciences*, 6(1), 256–267. https://doi.org/10.18502/kls.v6i1.8613
- Padilha, J. M. D. S. C., & Silva, R. P. (2020). Impacte da pandemia por covid-19 nos enfermeiros de reabilitação portugueses. *Revista Portuguesa de Enfermagem de Reabilitação*, 3(2), Art. 2. https://doi.org/10.33194/rper.2020.v3.s2.15.5842
- Pérez-Fuentes, M. del C., Molero-Jurado, M. del M., Gázquez-Linares, J. J., & Simón-Márquez, M. del M. (2018). Analysis of Burnout Predictors in Nursing: Risk and Protective Psychological Factors. *European Journal of Psychology Applied to Legal Context*, 11(1), 33–40. https://doi.org/10.5093/ejpalc2018a13
- Ribeiro, J. L. P., & Marques, T. M. P. F. dos S. (2009). A avaliação do stresse: A propósito de um estudo de adaptação da escala de percepção de stresse. *Psicologia, Saúde & Doenças, 10*(2), 237–248.
- Teixeira, A. L., Krause, T. M., Ghosh, L., Shahani, L., Machado-Vieira, R., Lane, S. D., Boerwinkle, E., & Soares, J. C. (2021). Analysis of COVID-19 Infection and Mortality Among Patients With Psychiatric Disorders, 2020. JAMA Network Open, 4(11), e2134969. https://doi.org/10.1001/jamanetworkopen.2021.34969



- Temsah, M.-H., Al-Sohime, F., Alamro, N., Al-Eyadhy, A., Al-Hasan, K., Jamal, A., Al-Maglouth, I., Aljamaan, F., Al Amri, M., Barry, M., Al-Subaie, S., & Somily, A. M. (2020). The psychological impact of COVID-19 pandemic on health care workers in a MERS-CoV endemic country. *Journal of Infection and Public Health*, 13(6), 877–882. https://doi.org/10.1016/j.jiph.2020.05.021
- Urzal, M.; Donas- Boto, I.; Moreira, M.; Nogueira, P. & Vian, J. (2021). Prevalência e Fatores associados a sintomas de Ansiedade, Depressão e Perturbação Pós- stress Traumático em Profissionais de Saúde durante a Pandemia por COVID-19. (2021). *RPSO - Revista Portuguesa de Saúde Ocupacional, 11*, 1.23. https://doi.org/10.31252/RPSO.26.06.2021
- Walton, M., Murray, E., & Christian, M. D. (2020). Mental health care for medical staff and affiliated healthcare workers during the COVID-19 pandemic. *European Heart Journal. Acute Cardiovascular Care*, 9(3), 241–247. https://doi.org/10.1177/2048872620922795
- Zakeri, M. A., Hossini Rafsanjanipoor, S. M., Zakeri, M., & Dehghan, M. (2021). The relationship between frontline nurses' psychosocial status, satisfaction with life and resilience during the prevalence of COVID-19 disease. *Nursing Open*, 8(4), 1829–1839. https://doi.org/10.1002/nop2.832
- Zerbini, G., Ebigbo, A., Reicherts, P., Kunz, M., & Messman, H. (2020). Psychosocial burden of healthcare professionals in times of COVID-19 a survey conducted at the University Hospital Augsburg. *GMS German Medical Science*, *18*, Doc05. https://doi.org/10.3205/000281
- Zhang, Y., & Ma, Z. F. (2020). Impact of the COVID-19 Pandemic on Mental Health and Quality of Life among Local Residents in Liaoning Province, China: A Cross-Sectional Study. International Journal of Environmental Research and Public Health, 17(7), 2381. https://doi.org/10.3390/ijerph17072381