

# The Influence of Pregestational Body Mass Index and Physical Activity Patterns on Maternal, Delivery, and Newborn Outcomes in a Sample of Portuguese Pregnant Women: A Retrospective Cohort Study

Diana Bernardo<sup>a, b</sup> Carlos Carvalho<sup>c</sup> Jorge Mota<sup>b, d</sup> Margarida Ferreira<sup>e, f</sup>  
Paula Clara Santos<sup>b, d, g, h</sup>

<sup>a</sup>KinesioLab Research Unit in Human Movement, Department of Physiotherapy, School of Health, Piaget Institute, Vila Nova de Gaia, Portugal; <sup>b</sup>Faculty of Sport, Research Centre in Physical Activity, Health and Leisure (CIAFEL), Porto, Portugal; <sup>c</sup>Department of Physiotherapy, Sword Health Technologies, Porto, Portugal; <sup>d</sup>Laboratory for Integrative and Translational Research in Population Health (ITR), FADEUP- Faculty of Sport, University of Porto, Porto, Portugal; <sup>e</sup>Physiotherapy Department, CESPU, Polytechnic Health Institute of the North, Gandra, Portugal; <sup>f</sup>Physical and Rehabilitation Medicine, Hospital Senhora da Oliveira, Guimaraes, Portugal; <sup>g</sup>Department of Physiotherapy ESS, Polytechnic of Porto, Porto, Portugal; <sup>h</sup>Center for Rehabilitation Research (CIR), School of Health, Polytechnic of Porto, Porto, Portugal

## Keywords

Physical activity · Pregnancy · Pregnant women · Body mass index · Obesity · Pre-obesity

## Abstract

**Introduction:** There is a linear association between pre-gestational body mass index (BMI) and almost all adverse pregnancy outcomes. Pregnancy is “a window of opportunities” in terms of changing behavior and improving awareness of healthy living. The proper assessment of physical activity levels, during pregnancy, determines trends, health benefits, and their effects over time. This study aims to describe maternal physical activity levels, stratified by pregestational BMI, verify the accomplishment of physical activity recommendations in pregnant women, and correlate pregestational BMI and physical activity accomplishment with maternal, delivery, and neonatal parameters.

**Methods:** A retrospective cohort study was carried out with 103 pregnant women. Physical activity levels were evaluated using a questionnaire and accelerometry. Pregestational BMI was obtained through the Quetelet formula and used the American College of Sports Medicine’s guidelines were used to determine physical activity accomplishment levels. Continuous data were presented as mean and standard deviation and categorical data as numbers and percentages. The *F* test was used to examine the differences between groups. **Results:** Pregnant women in the sample spent 42.9% of their time on household activities, and for pregnant women with obesity, 91.5% of the time was spent on sedentary activities. Women with normative BMI had higher levels of moderate-intensity activities. Only 15.8% of participants with obesity reached the international recommendations for physical activity practice and women who accomplished physical activity recommendations gained less weight during pregnancy. A high percentage of

pre-obesity and obese pregnant women exceeded the recommendations for gestational weight gain and the gestational diabetes prevalence was higher in the obesity group ( $p = 0.03$  between groups). Regarding delivery and neonatal parameters (Apgar score 1st, Apgar score 5th, birth weight, length and head circumference), no statistical differences were found when adjusted to a gestational week at birth, between BMI ( $p = 0.58$ ;  $p = 0.18$ ;  $p = 0.60$ ;  $p = 0.34$ ;  $p = 0.34$ , respectively) or physical activity ( $p = 0.12$ ;  $p = 0.15$ ;  $p = 0.83$ ;  $p = 0.70$ ;  $p = 0.70$ , respectively) groups. **Conclusion:** Pregnant women with obesity, exhibit high levels of sedentary behavior, a high prevalence of gestational diabetes, and exceed recommended gestational weight gain. Healthcare professionals have a crucial role in promoting regular physical activity and lifestyle changes before and during pregnancy.

© 2023 The Author(s). Published by S. Karger AG, Basel on behalf of NOVA National School of Public Health

### A influência do Índice de Massa Corporal pré-gestacional e das características da atividade física nos parâmetros maternos, de parto e do recém-nascido numa amostra de grávidas portuguesas: estudo de coorte retrospectivo

#### Palavras Chave

Atividade física · Gravidez · Mulheres grávidas · IMC · Obesidade · Pré-obesidade

#### Resumo

**Introdução:** Existe uma associação linear entre o Índice de Massa Corporal (IMC) pré-gestacional e quase todos os desfechos adversos da gravidez. A gravidez é “uma janela de oportunidades” em termos de mudança de comportamento e melhoria da conscientização sobre uma vida saudável. A avaliação adequada dos níveis de Atividade Física, durante a gravidez, determina tendências, benefícios para a saúde e os seus efeitos ao longo do tempo. Este estudo tem como objetivo descrever as atividades e os níveis de atividade física materna, estratificados pelo IMC pré-gestacional, verificar o cumprimento das recomendações de atividade física em grávidas e correlacionar o IMC e a realização de atividade física com parâmetros maternos, de parto e neonatais.

**Métodos:** Foi realizado um estudo de coorte retrospectivo com 103 grávidas. O tipo e a intensidade da atividade física foram avaliados por questionário e acelerometria, o IMC pré-gestacional foi obtido pela fórmula de Quetelet e

utilizaram-se as diretrizes do *American College of Sport Medicine* para determinar os níveis de realização da atividade física. Os dados contínuos foram apresentados em média e desvio padrão, e os dados categóricos em números e percentagens. O *F test* e foi utilizado para examinar as diferenças entre os grupos. **Resultados:** As grávidas da amostra despenderam 42.9% do tempo em atividades domésticas e, as grávidas com obesidade, 91.5% do tempo em atividades sedentárias. Grávidas com IMC normativo apresentaram níveis mais elevados de atividades de intensidade moderada, 15.8% das participantes com obesidade atingiram as recomendações internacionais para a prática de atividade física e as mulheres que cumpriram as recomendações de atividade física ganharam menos peso durante a gravidez. Grande percentagem das grávidas com sobrepeso e obesidade não cumpriu com as recomendações para ganho de peso gestacional e a incidência de diabetes gestacional foi maior no grupo obesidade ( $p = 0.03$  entre os grupos). Relativamente aos parâmetros de parto e neonatais (apgar 1o, apgar 5o, peso tamanho e perímetro cefálico), não foram encontradas diferenças estatísticas, quando os scores foram ajustados à semana de gestacional, entre os grupos de IMC ( $p = 0.58$ ;  $p = 0.18$ ;  $p = 0.60$ ;  $p = 0.34$ ;  $p = 0.34$  respetivamente) ou atividade física ( $p = 0.12$ ;  $p = 0.15$ ;  $p = 0.83$ ;  $p = 0.70$ ;  $p = 0.70$  respetivamente). **Conclusão:** Grávidas com obesidade apresentam altos níveis de comportamento sedentário, alta prevalência de diabetes gestacional e excedem o ganho de peso gestacional recomendado. Os profissionais de saúde têm um papel crucial na promoção da atividade física regular e mudanças no estilo de vida antes e durante a gravidez.

© 2023 The Author(s). Published by S. Karger AG, Basel on behalf of NOVA National School of Public Health

#### Introduction

The attitude toward physical activity (PA) during pregnancy has changed over the last 50 years [1]. In the past, when women became pregnant, were promptly advised to reduce or even discontinue the practice of PA due to the risk of complications for the mother and fetus [2]. This was mainly due to sociocultural factors and a lack of scientific evidence to demonstrate safety in this practice [3].

Nowadays PA, defined as any voluntary bodily movement produced by musculoskeletal contraction that results in a substantial increase in caloric requirements over resting energy expenditure [4], is not

only considered safe for the mother and fetus but also improves important pregnancy outcomes [5, 6]. Studies show that PA is closely associated with reducing the risk of obesity, hypertension, diabetes [7–9], cardiovascular disease [10, 11], increased emotional well-being [12, 13], reducing the risk of postpartum depression [7, 8, 14, 15], preeclampsia [12], complications during childbirth [16], and risk of cesarean delivery [12].

There is a linear association between pregestational body mass index (BMI) and almost all adverse pregnancy outcomes [17, 18]. Pregnant women with an elevated BMI are at increased risk for the development of gestational diabetes mellitus, hypertensive disorders of pregnancy, and a need for unplanned cesarean deliveries. Their infants are also at increased risk for prematurity, stillbirth, macrosomia, birth trauma, respiratory distress, hypoglycemia, early postnatal infection, neonatal intensive care unit admission, and neonatal death [17].

Obesity in pregnant women, defined as pregestational BMI  $\geq 30$  kg/m<sup>2</sup>, is a significant public health challenge [19], worldwide, there are almost 15 million pregnant women with obesity [20]. In the USA alone, 29% of women had obesity before becoming pregnant [19].

The proportion of pregnant women of reproductive age is increasing worldwide [21]. In Portugal, 49.3% of the female population over 18 years old has excess weight or obesity [22], and a small study with 88 Portuguese pregnant women showed that 14.8% of them had obesity.

Pregnancy is a “window of opportunities” in terms of changing behavior and improving awareness of healthy living [23]. There is increasing interest in research in the general population about whether reducing time spent in sedentary behaviors has a beneficial effect on health [24]. Therefore, it became important to properly assess PA levels in different pregestational BMI pregnant women. PA evaluation determines trends, health benefits, and their effects over time [25]. The purpose of this study was to (1) describe maternal PA levels and activities, stratified by pregestational BMI; (2) verify the accomplishment of PA recommendations in pregnant women with normative weight, pre-obesity, or obesity; (3) correlate pregestational BMI and PA accomplishment with maternal, delivery, and neonatal parameters.

## Material and Methods

We conducted a retrospective cohort study at the University Hospital Center of São João (CHUSJ), Porto, at two different times. Women were recruited and assessed during medical appointments between July 2010 and May 2012, and May 2019 and August 2019.

We exclude pregnant women: (1) those under 18 years of age; (2) diagnosed with insulin-dependent diabetes, hypertension, and/or cardiac diseases; (3) multiple pregnancies; and (4) those unable to follow or understand the procedure or who did not provide informed consent to participate in research [26, 27]. A total of 264 pregnant women were invited to participate in the study and 54 refused. Of those who agreed, 29 were lost in follow-up. The final sample, which included women who participated in both moments, consisted of 103 pregnant women. Figure 1 shows the flow chart of participants. We examined the following maternal outcomes: PA levels, gestational weight gain (GWG), excessive gestational weight gain (EGWG), and gestational diabetes mellitus (GDM); the delivery outcomes were: the type of birth and gestational age of birth; the newborn outcomes were: the Apgar score (1st and 5th min), weight at born, body composition, and head circumference. The sociodemographic data were collected by a standardized questionnaire.

### *Pregnant Women Evaluation*

The pregnant women were recruited for our study during different gestational trimesters. PA evaluation was carried out once during the 1st, 2nd, or 3rd gestational trimester at the same time, we collected sociodemographic data. We followed up until delivery.

### *PA Measurement*


Proper assessments of PA are important for determining both trends and health benefits and their effects over time. PA can be evaluated using objective methods, such as accelerometers and/or subjectively based on questionnaires [28]. The PA was objectively assessed by accelerometry and subjectively assessed by questionnaire.

The accelerometers Actigraph GT3X<sup>®</sup> and wGT3X-BT<sup>®</sup> used in this study, monitored the acceleration of body segments and are considered a gold standard in objectivity and reliability (ICC: 0.661–0.806) measuring sure PA levels [29]. They have previously been used in pregnant women, including pregnant women with obesity, and showed good acceptability [26, 30]. Pregnant women wear the accelerometer on the waist for 7 consecutive days. Only those that record at least 480 miles daily, for a least 3 days were considered valid [29]. The accelerometry of different intensity levels was accessed by Freedson [27] with cut points, which have already been used in other studies with pregnant women [31, 32].

The Pregnancy Physical Activity Questionnaire (PPAQ) is a short-form questionnaire with 36 questions that aim to measure the type of activity and intensity of PA in pregnant women. It is a valid and reliable instrument ( $r = 0.09$ – $0.43$ ; ICC  $>0.78$ ) that assesses the duration, frequency and intensity of household/caregiving, occupational, sports/exercise, transport and inactivity activities during the trimester and provides a quantitative measure of PA intensity which is classified as sedentary ( $<1.5$  METs), light ( $1.5$ – $3.0$  METs), moderate ( $3.1$ – $6.0$  METs), or vigorous ( $>6.0$  METs). One MET corresponds to the metabolic equivalent of energy expended at rest [32]. It is a specific questionnaire for this population group and recommended in the literature to subjectively assess PA in pregnancy [33].

### *PA Recommendations*

The American College of Sports Medicine (ACSM) recommends that pregnant women practice at least 150 min of moderate/vigorous physical activity intensity per week throughout the entire pregnancy [34].



**Fig. 1.** Chart of exclusions and losses in the cohort.

#### Anthropometric Measures

The pregnant woman BMI's was calculated using the Quetelet formula, which is based on individual height and weight – BMI = weight (kg)/(height)<sup>2</sup> (m). The self-reported pregestational weight was confirmed in the Pregnant Health Bulletin, and the height was obtained through a fixed stadiometer (SECA 206). The weight was collected using a portable digital scale (Tantita InnerScan BC-545). According to the World Health Organization (WHO), the BMI can be categorized as underweight (BMI <18.4 kg/m<sup>2</sup>), normative weight (BMI 18.5–24.9 kg/m<sup>2</sup>), pre-obesity (BMI 25–29.9 kg/m<sup>2</sup>), and obesity (BMI ≥30 kg/m<sup>2</sup>) [35].

#### Gestational Weight Gain and Excessive Gestational Weight Gain

GWG was calculated as the difference between maternal weight at the last appointment (36–39 gestational weeks) and pregestational weight. The ideal GWG during pregnancy is defined by IOM (2009) and depends on pregestational BMI [36]. The adequate GWG in normative pregestational BMI women ranges between 11.5 and 16 kg; in pre-obesity, between 7 and 11.5 kg and in women with obesity, it should not exceed 9 kg in the entire pregnancy. Values over these thresholds are considered EGWG [37].

#### Gestational Diabetes Mellitus

The presence of GDM was assessed between 24 and 28 gestation weeks through the oral glucose tolerance test. Blood sampling for fasting glucose concentrations was taken after a 10-h overnight fast, and glucose tolerance was measured by a 2-h 75 mg per-oral glucose tolerance test. GDM was diagnosed as fasting glucose ≥126 mg/dL or 2 h concentration ≥200 mg/dL [38].

#### Birth and Offspring Outcomes

The birth weight cut-offs were established by WHO Child Growth Standards as low birth weight (≤10th percentile), normal birth weight (>10th percentile <90th), and macrosomia (≥90th percentile) [39]. For detailed analysis, the birth weight, length, and head circumferences Z-scores were predefined in the following percentiles: Z-score –4 to < –2 (0 percentile), Z-score –2 to < –1 (2nd percentile), Z-score –1 to < –0.5 (16th percentile), Z-score –0.5 to < 0 (31st percentile), Z-score 0 (50th percentile), Z-score 0.5 to < 1 (69th percentile), Z-score 1 to < 2 (84th percentile), Z-score 3–4 (100th percentile). Data were collected by consulting the CHUSJ patient management computer system and Newborn Child Health Bulletin.

#### Statistics

Continuous data were presented in mean and standard deviation unless otherwise specified. Categorical data were presented as numbers and percentages. The *F* test in analysis of variance (ANOVA) was used to examine differences in characteristics between groups, for continuous variables. When normality was not assumed, we used the Kruskal-Wallis test. The  $\chi^2$  test was used for categorical variables. The mother's age, the newborn's sex and the gestational week can influence maternal and offspring outcomes [40]. These potentially confounding variables were adjusted to gestational week at birth in the multivariate logistic regression models to calculate maternal GWG, and offspring weight, length, and head circumference.

Statistical analyses were performed using SPSS for Windows, Version 18 (IBM, Armonk, NY). *p* < 0.05 was considered to indicate a statistically significant relationship.

**Table 1.** Sociodemographic and obstetric characteristics at baseline

Age, years ( $n = 103$ ) <sup>a</sup>	30.31 ( $\pm 5.5$ )
Prepregnancy BMI ( $n = 103$ ) <sup>a</sup>	29.1 ( $\pm 6.88$ )
Normative, $n$ (%)	35 (34.0)
Pre-obesity, $n$ (%)	19 (18.4)
Obesity, $n$ (%)	49 (47.6)
Gestational trimester ( $n = 103$ )	
1st trimester, $n$ (%)	40 (38.8)
2nd trimester, $n$ (%)	34 (33.0)
3rd trimester, $n$ (%)	29 (28.2)
Education level ( $n = 98$ )	
Mandatory or less, $n$ (%)	32 (32.6)
Secondary, $n$ (%)	35 (35.7)
College/university, $n$ (%)	31 (31.6)
Professional status ( $n = 98$ )	
Employed/student, $n$ (%)	78 (79.6)
Unemployed, $n$ (%)	20 (20.4)
Marital status ( $n = 98$ )	
Married/cohabitated, $n$ (%)	76 (77.6)
Single/divorced, $n$ (%)	22 (22.6)
Number of gestations ( $n = 103$ )	
Primigest, $n$ (%)	31 (30.1)
Multigest, $n$ (%)	72 (69.9)

BMI, body mass index; SD, standard deviation. <sup>a</sup>Data are mean  $\pm$  standard deviation.

## Results

### Sociodemographic Characteristics

The mean age of women was 30.31 years ( $\pm 5.5$ ). Among 103 participants, 35 (34.0%) had a normative pre-pregnant BMI (range 20.8–24.5 kg/m<sup>2</sup>), 19 (18.4%) pre-obesity, and 49 (47.6%) obesity (range 30.4–47.5 kg/m<sup>2</sup>). At the recruitment moment, 40 pregnant women (38.8%) were in the first gestational trimester, 34 (33.0%) were in the second gestational trimester, and 29 (28.2%) were in the third. Thirty-one (31.6%) pregnant women had a high education level, 78 (79.6%) were employed, and 76 (77.6%) were married or cohabitated. The sociodemographic characteristics are shown in Table 1.

### PA Assessed by Self-Reported Questionnaire

Table 2 represents maternal PA levels measured by PPAQ. The data show that pregnant women spent most of their time (42.9%) on household/caregiving activities and in occupational activities (36.8%). The results also show a very small percentage of energy expenditure in sports/exercise activities (2.5%). Regarding PA intensity measured by PPAQ, 42.6% of total activities were in light intensities, and just 0.54% of activities were spent in activities above 6METs. When the data were stratified by pregestational BMI, the results were similar between groups.

### PA Assessed by Accelerometry

The accelerometry results (Table 3) indicate that pregnant women in the sample spent 96.5% of the time in activities below 3 METs of expended energy (light or below) and 3.1% of the time in moderate activities (between 3.1 and 6.0 METs). When data were stratified by pregestational BMI, we observed that women with normative BMI had higher levels in moderate activities, compared with pre-obesity and obesity BMI groups, with statistical differences between them ( $p = 0.006$ ). Pregnant women with obesity spent 91.5% of their time in sedentary activities. Time spent in light-intensity activities in normative, pre-obesity, and obesity BMI groups were 15.1%, 7.8%, and 5.9%, respectively ( $p < 0.001$ ). Regarding vigorous activities, pregnant women with normative pregestational BMI spent 1.2% of the time in activities over 6METs, and pregnant women with pre-obesity and obesity did not spend any time on activities of this intensity ( $p < 0.001$ ).

### International PA Recommendations

44.7% of participants reached the international recommendations for PA practice. In pregestational BMI groups, we found a statistically significant difference ( $p = 0.02$ ). 51.4% cent of normative weight and 51.0% of women with obesity achieved the recommendations, while only 15.8% of pregnant women with pre-obesity reached the threshold.

### Maternal, Delivery, and Neonatal Parameters

The results (Table 4) showed that pregnant women gained a mean of 13.09 kg ( $\pm 6.41$ ) during the entire pregnancy. Statistical differences were found ( $p = 0.03$ ) between women who meet PA recommendations (11.21 kg [ $\pm 6.21$ ]) and women who did not meet the criteria (14.61 kg [ $\pm 6.20$ ]). Regarding pregestational BMI (Table 5), pregnant women with obesity gained lower weight [11.13 kg ( $\pm 6.93$ )] compared to normative pregnant women (15.57 kg [ $\pm 5.87$ ]) with a statistical difference between groups ( $p = 0.01$ ). However, the data showed that 68.4% of pre-obesity and 65.3% of obese pregnant women exceed the IOM recommendation for GWG.

GDM was present in 14.6% of participants, with a statistical difference between pregestational BMI groups ( $p = 0.03$ ). The obesity group had a GDM prevalence of 24.5% compared to 5.7% in the normative weight group.

39.4 weeks ( $\pm 1.63$ ) was the mean of the gestational age at birth for the total sample. 39.8% of total participants had a cesarean delivery, with the highest percentage in pregnant women who did not meet the international recommendations

**Table 2.** Physical activity levels assessed by PPAQ

Intensity	Total sample (n = 103)		Normative BMI (n = 35)		Pre-obesity BMI (n = 19)		Obesity BMI (n = 49)		p value <sup>1</sup>
	(MET.h/wk) <sup>a</sup>	% <sup>b</sup>	(MET.h/wk) <sup>a</sup>	% <sup>b</sup>	(MET.h/wk) <sup>a</sup>	% <sup>b</sup>	(MET.h/wk) <sup>a</sup>	% <sup>b</sup>	
Total	264.81 (±152.06)	100	239.95 (±114.50)	100	266.42 (±186.74)	100	288.95 (±165.89)	100	0.87
Sedentary	50.75 (±32.52)	19.2	50.41 (±32.01)	20.6	39.88 (±28.03)	15.0	57.14 (±33.99)	19.8	0.17
Light	112.80 (±56.56)	42.6	109.51 (±55.11)	45.6	111.15 (±61.15)	41.7	116.86 (±56.70)	40.4	0.96
Moderate	99.83 (±110.41)	37.7	79.39 (±74.69)	33.1	114.36 (±143.95)	42.9	113.56 (±121.31)	39.3	0.69
Vigorous	1.43 (±3.36)	0.54	1.64 (±3.67)	0.7	1.03 (±2.80)	0.4	1.40 (±3.32)	0.5	0.52
Type									
Household/caregiving	113.61 (±77.52)	42.9	106.14 (±74.54)	44.2	112.34 (±74.54)	42.2	121.68 (±79.55)	42.1	0.98
Occupational	97.51 (±111.97)	36.8	84.75 (±80.84)	35.3	104.81 (±80.93)	39.3	106.91 (±130.06)	37.0	0.97
Sports/exercise	6.76 (±7.88)	2.5	7.20 (±9.04)	3.0	5.80 (±5.82)	2.2	6.77 (±7.55)	2.3	0.72
Transportation	22.68 (±19.03)	8.6	19.92 (±16.24)	8.3	22.71 (±20.64)	8.5	25.43 (±20.75)	8.8	0.65
Inactivity	24.26 (±18.98)	9.2	21.64 (±16.55)	9.0	20.75 (±14.93)	7.8	28.16 (±22.34)	9.7	0.36

<sup>a</sup>Data are mean ± standard deviation. <sup>b</sup>Used mean to calculate percentages. <sup>1</sup>Analysis by Kruskal-Wallis test between groups.

**Table 3.** Physical activity levels assessed by accelerometry

Intensity	Total sample (n = 103)		Normative BMI (n = 35)		Pre-obesity BMI (n = 19)		Obesity BMI (n = 49)		p value <sup>1</sup>
	min/day <sup>a</sup>	% <sup>b</sup>	min/day <sup>a</sup>	% <sup>b</sup>	min/day <sup>a</sup>	% <sup>b</sup>	min/day <sup>a</sup>	% <sup>b</sup>	
Total	983.20 (±281.89)	100	818.63 (±235.37)	100	1,126.81 (±266.82)	100	1,139.30 (±198.46)	100	<0.001
Sedentary	851.10 (±295.47)	86.6	650.50 (±246.75)	79.5	1,016.63 (±239.89)	90.2	1,042.07 (±186.11)	91.5	<0.001
Light	97.67 (±49.40)	9.9	123.58 (±49.23)	15.1	87.82 (±36.73)	7.8	66.75 (±34.34)	5.9	<0.001
Moderate	30.55 (±16.77)	3.1	34.54 (±16.42)	4.2	22.07 (±11.50)	2.0	29.58 (±18.14)	2.6	0.006
Vigorous	3.88 (±18.84)	0.4	10.01 (±31.68)	1.2	0.29 (±0.22)	0.0	0.90 (±1.22)	0.0	<0.001
	Total sample (n = 103)		Normative BMI (n = 35)		Pre-obesity BMI (n = 19)		Obesity BMI (n = 49)		p value <sup>1</sup>
	150 min/wk MVPA, n (%)		150 min/wk MVPA, n (%)		150 min/wk MVPA, n (%)		150 min/wk MVPA, n (%)		
<i>International PA recommendations accomplishment</i>	46 (44.7)		18 (51.4)		3 (15.8)		25 (51.0)		0.02

BMI, body mass index; MVPA, moderate/vigorous physical activity; min/wk, minutes/week. <sup>a</sup>Data are mean ± standard deviation. <sup>b</sup>Used mean to calculate percentages. <sup>1</sup>Analysis by Kruskal-Wallis test between groups.

for PA practice (43.9%) and women with a pregestational BMI categorized as pre-obesity (52.6%). However, no statistical differences were found between the groups.

The neonatal parameters showed an Apgar score at the 1st and 5th minute of 9 and 10, respectively, the birth weight mean was 3303.79 g (±423.79), the length mean was 49.66 cm (±2.38) and the head circumference mean was 38.23 cm (±32.84). No statistical differences were found between those who met/not met PA practice recommendations or between pregestational BMI groups ( $p > 0.05$ ). No statistical differences were found when values of potentially confounding variables (mother's age, newborn sex) were adjusted.

## Discussion

This study aimed to describe maternal PA levels and activities stratified by pregestational BMI, verify the achievement of PA recommendations, and correlate pregestational BMI and PA achievement with maternal, delivery, and neonatal parameters. Our key findings suggest that household/caregiving and occupational activities constituted the largest proportion of the total activity levels in pregnant women. Pregnant women with obesity had higher levels of sedentary behavior, higher prevalence of GDM, and exceeded the international recommendations for GWG. The prevalence of women

**Table 4.** Relationship between maternal, delivery, and neonatal parameters to the international PA recommendations accomplishment group

	Total sample (n = 103)	International PA recommendations		
		accomplished (n = 46)	not accomplished (n = 57)	p value
<b>Maternal parameters</b>				
GWG, kg <sup>a</sup>	13.09 (±6.41)	11.21 (±6.21)	14.61 (±6.20)	<b>0.03<sup>3</sup></b>
GWG adjusted to gestational week at birth	-0.00 (±0.16)	-0.01 (±0.19)	-0.00 (±0.12)	<b>0.04<sup>3</sup></b>
Exceed GWG, n (%)	56 (54.4)	24 (52.7)	22 (38.6)	<b>0.07<sup>1</sup></b>
Gestational diabetes mellitus, n (%)	15 (14.6)	7 (15.2)	8 (14.0)	<b>0.97<sup>1</sup></b>
<b>Delivery parameters</b>				
Gestational age on birth, weeks <sup>a</sup>	39.45 (±1.63)	39.33 (±2.03)	39.54 (1.23)	<b>0.74<sup>3</sup></b>
Type of birth				
Vaginal, n (%)	34 (33.0)	13 (28.2)	21 (36.8)	<b>0.20<sup>1</sup></b>
Instrumental vaginal, n (%)	28 (27.2)	17 (37.0)	11 (19.3)	
Cesarian, n (%)	41 (39.8)	16 (34.8)	25 (43.9)	
<b>Neonatal parameters</b>				
Apgar score 1st minute <sup>b</sup>	8.56 (±1.14)	8.35 (±1.54)	8.74 (±0.64)	<b>0.12<sup>3</sup></b>
Apgar score 5th minute <sup>b</sup>	9.61 (±1.17)	9.41 (±1.64)	9.77 (±0.54)	<b>0.15<sup>3</sup></b>
Birth weight, g <sup>a</sup>	3,303.79 (±423.79)	3,290.98 (±403.88)	3,314.12 (±449.80)	<b>0.78<sup>2</sup></b>
Birth weight cut-offs				
Low birth weight, n (%)	34 (33)	17 (37)	17 (29.8)	<b>0.36<sup>1</sup></b>
Normal weight, n (%)	67 (65)	29 (63)	38 (66.7)	
Macrosomia, n (%)	2 (2)	0 (0)	2 (3.5)	
Birth weight z-score adjusted to gestational week at birth, g <sup>a</sup>	0.01 (±0.37)	0.00 (±0.44)	0.01 (±0.28)	<b>0.83<sup>2</sup></b>
Length, cm <sup>a</sup>	49.66 (±2.38)	49.58 (±2.89)	49.72 (±1.90)	<b>0.42<sup>3</sup></b>
Length z-score adjusted to gestational week at birth, cm <sup>a</sup>	-0.00 (±0.25)	-0.01 (±0.30)	-0.00 (±0.19)	<b>0.70<sup>3</sup></b>
Head circumference, cm <sup>a</sup>	38.23 (±32.84)	42.56 (±48.86)	34.73 (±4.20)	<b>0.52<sup>3</sup></b>
Head circumference z-score adjusted to gestational week at birth, cm <sup>a</sup>	0.25 (±0.12)	-0.00 (±0.15)	-0.00 (±0.09)	<b>0.70<sup>3</sup></b>

<sup>a</sup>Data are mean ± standard deviation. <sup>b</sup>Data are median (interquartile range). <sup>1</sup>Analysis by  $\chi^2$  test between groups. <sup>2</sup>Analysis by ANOVA test between groups. <sup>3</sup>Analysis by Kruskal-Wallis test between groups.

who reached the recommended level of PA is 44.7%; however, only 15.8% of pregnant women with pre-obesity engaged in activities above the threshold.

In this study, we found that, independently of pre-gestational BMI, more than 79% of PA time was engaged in household/caregiving and occupational activities. These results support other authors who have suggested that domestic and obligatory activities (work-related) were those that contributed the most to energy costs and sports/exercise-related activities were not important contributors to energy expenditure during pregnancy [30, 41].

Regarding the PA intensity, measured by self-report questionnaire (PPAQ), pregnant women of the sample spent 19.2% of their time on sedentary activities, and we did not find differences between pregestational BMI groups. However, when objectively assessed by accel-

erometry, the results showed that pre-obesity and obesity groups spent more than 90% of their time in sedentary activities. In Chandonnet et al. [26] study, the explanation for the differences between assessment instruments might be that pregnant women with obesity spend more time in light-intensity activities, but perceive it as moderate or vigorous.

The findings of this study indicate that pregestational BMI was inversely correlated with PA levels. These results are in line with other works, a 2017 systematic review demonstrated that pregnant women spend at least half of their time in sedentary activities [24] and women with pre-obesity or obesity before gestation were at increased risk of sedentary behavior during pregnancy [42]. Vigorous activities are extremely low in pregnant women [43] and women with obesity in general [44].

**Table 5.** Relationship between maternal, delivery, and neonatal parameters to the prepregnancy body mass index group

	BMI			p value
	normative BMI (n = 35)	pre-obesity BMI (n = 19)	obesity BMI (n = 49)	
<b>Maternal parameters</b>				
GWG, kg <sup>a</sup>	15.57 (±5.87)	13.58 (±4.09)	11.13 (±6.93)	<b>0.01<sup>3</sup></b>
GWG adjusted to gestational week at birth	-0.01 (±0.21)	0.03 (±0.13)	-0.01 (±0.12)	<b>0.03<sup>3</sup></b>
Exceed GWG, n (%)	11 (31.4)	13 (68.4)	32 (65.3)	<b>0.04<sup>1</sup></b>
Gestational diabetes mellitus, n (%)	2 (5.7)	1 (5.3)	12 (24.5)	<b>0.03<sup>1</sup></b>
<b>Delivery parameters</b>				
Gestational age on birth, weeks <sup>a</sup>	6.93 (±2.18)	39.79 (±1.36)	39.39 (±1.24)	0.34 <sup>3</sup>
Type of birth				
Vaginal, n (%)	12 (34.4)	6 (31.6)	16 (32.7)	0.64 <sup>1</sup>
Instrumental vaginal, n (%)	12 (34.3)	3 (15.8)	13 (26.5)	
Cesarian, n (%)	11 (31.4)	10 (52.6)	20 (40.8)	
<b>Neonatal parameters</b>				
Apgar score 1st minute <sup>b</sup>	8.77 (±0.49)	8.74 (±0.56)	8.35 (±1.55)	0.58 <sup>3</sup>
Apgar score 5th minute <sup>b</sup>	9.83 (±0.38)	9.84 (±0.37)	9.37 (±1.63)	0.18 <sup>3</sup>
Birth weight, g <sup>a</sup>	3,322.57 (±413.03)	3,378.68 (±315.95)	3,261.33 (±468.04)	0.56 <sup>2</sup>
Birth weight cut-offs				
Low birth weight, n (%)	12 (34.2)	5 (26.3)	17 (34.7)	0.89 <sup>1</sup>
Normal weight, n (%)	22 (62.9)	14 (73.7)	31 (63.3)	
Macrosomia, n (%)	1 (2.9)	0 (0)	1 (2.0)	
Birth weight z-score adjusted to gestational week at birth, g <sup>a</sup>	-0.02 (±0.49)	0.08 (±0.31)	-0.01 (±0.28)	0.60 <sup>2</sup>
Length, cm <sup>a</sup>	49.79 (±3.30)	49.58 (±0.96)	49.60 (±1.98)	0.37 <sup>3</sup>
Length z-score adjusted to gestational week at birth, cm <sup>a</sup>	-0.02 (±0.33)	0.05 (±0.20)	-0.01 (±0.19)	0.34 <sup>3</sup>
Head circumference, cm <sup>a</sup>	44.39 (±55.87)	35.66 (±6.97)	34.82 (±4.49)	0.42 <sup>3</sup>
Head circumference z-score adjusted to gestational week at birth, cm <sup>a</sup>	-0.01 (±0.16)	0.02 (±0.10)	0.09 (±0.00)	0.34 <sup>3</sup>

<sup>a</sup>Data are mean ± standard deviation. <sup>b</sup>Data are median (interquartile range). <sup>1</sup>Analysis by  $\chi^2$  test between groups; <sup>2</sup>Analysis by ANOVA test between groups. <sup>3</sup>Analysis by Kruskal-Wallis test between groups.

The regular practice of PA during pregnancy has shown multiple benefits. However notably, more than 50% of pregnant women in the sample failed to meet the recommendations of 150 weekly minutes of moderate/vigorous physical activity set in 2018 [34]. Nonetheless, results indicate that the rate of adherence to PA guidelines observed in this study is in line with Center for Disease Control (2018) [45] for the general population. When compared to other authors, we found that pregnant women in the sample had a higher rate of adherence to PA recommendations [46, 47]. This might be due to the fact that during pregnancy, women are in permanent contact with healthcare providers, and they encourage the regular practice of PA. Providers should be facilitators in promoting PA and lifestyle changes before and during pregnancy, especially in women with obesity who are typically more sedentary [48].

Pregnant women in the study gained 13.09 kg (±6.41) during pregnancy. The results have shown that women that did not achieve the PA recommendations gained more than 3.50 kg (mean) compared to those who meet the criteria. As occurred in other studies, PA and exercise seem to have a positive effect on GWG management [46, 49, 50]. When analyzing pregestational BMI groups, we verified that although the normative weight group gained more weight during pregnancy, more than 60% of pregnant women in pre-obesity and obesity groups exceeded the IOM recommendations for GWG. These findings corroborate other studies that demonstrate that a high percentage of pregnant women with pre-obesity or obesity exceed the IOM recommendations for GWG [50–52]. Healthcare providers to have a fundamental role in pregnant women's health. They should encourage pregnant women to remain active during pregnancy and adopt adjusted and healthy diets.



Our study showed that there is a solid association between maternal obesity and GDM, which is in line with other studies [54–56]. We did not find differences between groups that achieved or did not achieve PA recommendations. However, our results indicate that the normative weight group, which had higher levels of PA during pregnancy, was associated with a significantly lower risk of developing GDM, similar results were found in other studies [6–9, 57–60].

We also observed that 33.0% of women in this study had a vaginal delivery, 27.2% instrumental vaginal, and 39.8% a cesarean section delivery. The data between groups have shown no statistical differences. However, a higher rate of cesarean section delivery was found in pre-obesity and obesity groups. The high rates of cesarean delivery were suggested to be due to a possible link between increased cholesterol deposits in the myometrium of women with obesity, affecting contractions, an increase in maternal soft tissue inside the pelvis narrowing the birth canal and increasing difficulty in births [54]. Although our study has shown no statistical difference between pregnant who achieved and did not achieve PA recommendations, results from other studies support the benefits of PA, particularly in terms of reducing unplanned cesarean delivery [46, 61, 62].

Fetuses of obese gravidas are at an increased risk of macrosomia and impaired growth and preterm [17, 55]. Contrary to expectations, in our sample, we found no differences in birth weight cut-offs, length and head circumference, gestational age at birth and Apgar scores between pregestational BMI and achieve/not achieve PA recommendations groups.

Nevertheless, the findings of this study might present certain limitations. First, we only evaluate women in one gestational trimester. Due to that, we cannot verify PA changes during the entire pregnancy. Second, questionnaires are simple tools and easy to apply in research, however, self-reported measures can lead to possible classification errors [30]. This concern was minimized through the use of a developed and validated questionnaire for the pregnancy population [28, 32, 63] and with the use of accelerometry which evaluates objectively the PA levels. Future studies with tailored PA and exercises programs, specifically in pregnant women with pre-obesity and obesity, are needed to explore the impact of these factors on the health and quality of life of mother and offspring.

## Conclusion

Pregnant women with obesity, in this sample, exhibit high levels of sedentary behavior, a high prevalence of gestational diabetes, and exceed recommended GWG.

Healthcare professionals have a crucial role in promoting regular PA and healthy lifestyle changes before and during pregnancy.

In the same way that nutrition programs for pregnant women have been effective, physical exercise programs conducted in person or at home-based, by healthcare providers could be a useful method for increasing PA during pregnancy. This is especially true for populations with comorbidities such as pregnant women with obesity.

## Statement of Ethics

The Institutional Review Board of CHUSJ approved the original studies (n° 165/19) and reference number (09988). The use of the PPAQ questionnaire for this study was authorized by the author of the validation and adaptation for the Portuguese language and population. The study was conducted following the World Medical Association Helsinki Declaration for Human Studies. All participants provided written informed consent.

## Conflict of Interest Statement

The authors also declare that have no conflicts of interest to declare.

## Funding Sources

This work is supported by National Funds by FCT – Portuguese Foundation for Science and Technology, under the projects UI/BD/151206/2021 and CIAFEL: FCTUID/DTP/00617/2020; ITR: LA/P/0064/2020.

## Author Contributions

The conception, study design, and data acquisition were Diana Bernardo and Paula Clara Santos responsibility. Diana Bernardo and Carlos Carvalho analyzed and interpreted data and drafted the manuscript. The critical review regarding important intellectual content was carried out by Margarida Ferreira, Jorge Mota, and Paula Clara Santos. All authors approved the final version of the manuscript.

## Data Availability Statement

All data generated or analyzed during this study are included in this article. Further inquiries can be directed to the corresponding author.

## References

- 1 Newton ER, May L. Adaptation of maternal-fetal physiology to exercise in pregnancy: the basis of guidelines for physical activity in pregnancy. *Clin Med Insights Womens Heal*. 2017;10:1179562X1769322.
- 2 Briend A. Maternal physical activity, birth weight and perinatal mortality. *Med Hypotheses*. 1980;6(11):1157–70.
- 3 Downs DS, Chasan-Taber L, Evenson KR, Leiferman J, Yeo S. Physical activity and pregnancy: past and present evidence and future recommendations. *Res Q Exerc Sport*. 2012;83(4):485–502.
- 4 Caspersen CJ, Powell KE, Christenson G. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep*. 1985; 100(2):126–31.
- 5 Warburton DER, Bredin SSD. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr Opin Cardiol*. 2017;32(5):541–56.
- 6 Yu Y, Xie R, Shen C, Shu L. Effect of exercise during pregnancy to prevent gestational diabetes mellitus: a systematic review and meta-analysis. *J Matern Neonatal Med*. 2018; 31(12):1632–7.
- 7 Hayes L, McParlin C, Kinnunen TI, Poston L, Robson SC, Bell R, et al. Change in level of physical activity during pregnancy in obese women: findings from the UPBEAT pilot trial. *BMC Pregnancy Childbirth*. 2015; 15:e52.
- 8 Lauer EE, Jackson AW, Martin SB, Morrow JR. Meeting USDHHS physical activity guidelines and health outcomes. *Int J Exerc Sci*. 2017;10(1):121–7.
- 9 Zheng J, Wang H, Ren M. Influence of exercise intervention on gestational diabetes mellitus: a systematic review and meta-analysis. *J Endocrinol Invest*. 2017;40(10):1027–33.
- 10 Harrison CL, Thompson RG, Teede HJ, Lombard CB. Measuring physical activity during pregnancy. *West J Nurs Res*. 2011; 8(1):19–34.
- 11 Nascimento SL, Surita FG, Cecatti JG. Physical exercise during pregnancy: a systematic review. *Curr Opin Obstet Gynecol*. 2012;24(6):387–94.
- 12 ACOG. Committee. Opinion number 650: physical activity an exercise during pregnancy and the postpartum period. *Obstet Gynecol*. 2015;126(6):135–42.
- 13 Santo EC, Forbes PW, Oken E, Belfort MB. Determinants of physical activity frequency and provider advice during pregnancy. *BMC Pregnancy Childbirth*. 2017;17(1):e286.
- 14 Daley AJ, Foster L, Long G, Palmer C, Robinson O, Walmsley H, et al. The effectiveness of exercise for the prevention and treatment of antenatal depression: systematic review with meta-analysis. *BJOG An Int J Obstet Gynaecol*. 2015;122(1):57–62.
- 15 Mourady D, Richa S, Karam R, Papazian T, Hajj Moussa F, El Osta N, et al. Associations between quality of life, physical activity, worry, depression and insomnia: a cross-sectional designed study in healthy pregnant women. *PLoS One*. 2017;12(5): e0178181.
- 16 Foxcroft KF, Rowlands IJ, Byrne NM, McIntyre HD, Callaway LK; BAMBINO group. Exercise in obese pregnant women: the role of social factors, lifestyle and pregnancy symptoms. *BMC Pregnancy Childbirth*. 2011;11:4.
- 17 D'Souza R, Horyn I, Pavalagantharajah S, Zaffar N, Jacob CE. Maternal body mass index and pregnancy outcomes: a systematic review and metaanalysis. *Am J Obstet Gynecol MFM*. 2019;1(4):e100041.
- 18 Vinter CA, Nøhr E. Targeting pregnancy as a time to treat obesity. *Expert Rev Endocrinol Metab*. 2015;10(5):491–7.
- 19 Driscoll AK, Gregory ECW. Increases in prepregnancy obesity: United States, 2016–2019. *NCHS Data Brief*. 2020;392:1–8.
- 20 Chen C, Xu X, Yan Y. Estimated global overweight and obesity burden in pregnant women based on panel data model. *PLoS One*. 2018;13(8):e0202183–14.
- 21 OECD. *Health at a glance 2019: OECD indicators*. Paris: Organisation for Economic Co-operation and Development; 2019.
- 22 Projeção INE. *Da população residente com 18 e mais anos com excesso de peso ou obesidade (%) por Local de residência (NUTS - 2013), Sexo e tipologia de áreas urbanas: quinzenal: inquérito Nacional de Saúde: série 2014*. Lisboa: Instituto Nacional de Estatística; 2020. Available from: [https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine\\_indicadores&indOcorrCod=0010212&con texto=bd&selTab=tab2](https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&indOcorrCod=0010212&con texto=bd&selTab=tab2) (accessed 2022.01.12).
- 23 Poon LC, McIntyre HD, Hyett JA, da Fonseca EB, Hod M; FIGO Pregnancy and NCD Committee. The first-trimester of pregnancy: a window of opportunity for prediction and prevention of pregnancy complications and future life. *Diabetes Res Clin Pract*. 2018;145: 20–30.
- 24 Fazzi C, Saunders DH, Linton K, Norman JE, Reynolds RM. Sedentary behaviours during pregnancy: a systematic review. *Int J Behav Nutr Phys Act*. 2017;14(1):e32.
- 25 van Poppel MNM, Chinapaw MJM, Mokkink LB, van Mechelen W, Terwee CB. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sport Med*. 2010;40(7):565–600.
- 26 Chandonnet N, Saey D, Almérás N, Marc I. French pregnancy physical activity questionnaire compared with an accelerometer cut point to classify physical activity among pregnant obese women. *PLoS One*. 2012;7(6): e38818.
- 27 Chasan-Taber L, Schmidt MD, Pekow P, Sternfeld B, Manson JA, Markenson G. Correlates of physical activity in pregnancy among latina women. *Matern Child Health J*. 2007;11(4):353–63.
- 28 Bernardo D, Carvalho C, Leirós-Rodríguez R, Mota J, Santos PC. Comparison of the Portuguese Version of the Pregnancy Physical Activity Questionnaire (PPAQ) with Accelerometry for classifying physical activity among pregnant women with obesity. *Int J Environ Res Public Health*. 2023;20(2): 929–11.
- 29 Migueles JH, Cadenas-Sanchez C, Ekelund U, Delisle Nyström C, Mora-Gonzalez J, Löf M, et al. Accelerometer data collection and processing criteria to assess physical activity and other outcomes: a systematic review and practical considerations. *Sport Med*. 2017; 47(9):1821–45.
- 30 Santos PC, Abreu S, Moreira C, Santos R, Ferreira M, Alves O, et al. Physical activity patterns during pregnancy in a sample of Portuguese women: a longitudinal prospective study. *Iran Red Crescent Med J*. 2016; 18(3):e22455.
- 31 Freedson PS, Melanson E, Sirard J. Calibration of the computer science and applications, inc. Accelerometer. *Med Sci Sports Exerc*. 1998;30(5):777–81.
- 32 Chasan-Taber L, Schmidt MD, Roberts DE, Hosmer D, Markenson G, Freedson PS. Development and validation of a pregnancy physical activity questionnaire. *Med Sci Sports Exerc*. 2004;36(10):1750–60.
- 33 Sattler MC, Jaunig J, Watson ED, van Poppel MNM, Mokkink LB, Terwee CB, et al. Physical activity questionnaires for pregnancy: a systematic review of measurement properties. *Sport Med*. 2018; 48(10):2317–46.
- 34 Riebe D, Nobel M, Millholen A, editors 10th ed. *ACSM guideline for exercise testing and prescription*. Alphen aan den Rijn. the Netherlands: Wolters Kluwer Health; 2018.
- 35 WHO A. *Healthy lifestyle: WHO recommendations*; 2023. Available from: <https://www.who.int/europe/news-room/fact-sheets/item/a-healthy-lifestyle—who-recommendations> (accessed 2023.05.19).
- 36 Rasmussen KM, Yaktine AL, editors. *Weight gain during pregnancy: reexamining the guidelines*. Washington, DC: Institute of Medicine. National research council committee to reexamine IOM pregnancy weight guidelines. National Academies Press; 2009.
- 37 Rasmussen KM, Catalano PM, Yaktine AL. New guidelines for weight gain during pregnancy: what obstetrician/gynecologists should know. *Curr Opin Obstet Gynecol*. 2009;21(6):521–6.
- 38 Senat MV, Deruelle P. Gestational diabetes mellitus. *Diabetes Care*. 2016;44(4):244–7.

- 39 WHO. WHO child growth standards: book review. *Dev Med Child Neurol*. 2009;51(12):1002.
- 40 Calais-Ferreira L, Barreto ME, Mendonça E, Dite GS, Hickey M, Ferreira PH, et al. Birthweight, gestational age and familial confounding in sex differences in infant mortality: a matched co-twin control study of Brazilian male-female twin pairs identified by population data linkage. *Int J Epidemiol*. 2022;51(5):1502–10.
- 41 Cohen TR, Plourde H, Koski KG. Use of the Pregnancy Physical Activity Questionnaire (PPAQ) to identify behaviours associated with appropriate gestational weight gain during pregnancy. *J Phys Act Heal*. 2013;10(7):1000–7.
- 42 Andersen MB, Ostenfeld EB, Fuglsang J, Møller M, Daugaard M, Ovesen PG. Maternal prepregnancy body mass index and physical activity during pregnancy assessed by Accelerometer. *Am J Obstet Gynecol MFM*. 2020;2(4):e100182.
- 43 Rousham EK, Clarke PE, Gross H. Significant changes in physical activity among pregnant women in the UK as assessed by Accelerometry and self-reported activity. *Eur J Clin Nutr*. 2006;60(3):393–400.
- 44 Bond DS, Thomas JG, O’Leary KC, Lipton RB, Peterlin BL, Roth J, et al. Objectively measured physical activity in obese women with and without migraine. *Cephalalgia*. 2015;35(10):886–93.
- 45 Norris T, Schiller JS, Clarke TC; National Health Interview Survey Early Release Program. *Early release of selected estimates based on data from the 2018 National Health Interview Survey*. Washington, DC: National Center for Health Statistics. Centers for Disease Control and Prevention. Department of Health and Human Services; 2018.
- 46 Meander L, Lindqvist M, Mogren I, Sandlund J, West CE, Domellöf M. Physical activity and sedentary time during pregnancy and associations with maternal and fetal health outcomes: an epidemiological study. *BMC Pregnancy Childbirth*. 2021;21(1):e166.
- 47 Donofry SD, Germeroth LJ, Kolko Conlon RP, Venditti EM, Levine MD. Correlates of physical activity engagement among pregnant women with overweight and obesity. *Womens Heal Issues*. 2020;30(5):393–400.
- 48 Hsiung Y, Lee CF, Chi LK, Huang JP. “Moving for my baby!”: motivators and perceived barriers to facilitate readiness for physical activity during pregnancy among obese and overweight women of urban areas in Northern Taiwan. *Int J Environ Res Public Health*. 2021;18(10):e5275.
- 49 Muktabhant B, Lawrie TA, Lumbiganon P, Laopaiboon M. Diet or exercise, or both, for preventing excessive weight gain in pregnancy. *Cochrane Database Syst Rev*. 2015; 2015(6):CD007145.
- 50 Wang J, Wen D, Liu X, Liu Y. Impact of exercise on maternal gestational weight gain: an updated meta-analysis of randomized controlled trials. *Med*. 2019;98(27):e16199.
- 51 Ruiz JR, Perales M, Pelaez M, Lopez C, Lucia A, Barakat R. Supervised exercise-based intervention to prevent excessive gestational weight gain: a randomized controlled trial. *Mayo Clin Proc*. 2013;88(12):1388–97.
- 52 Bisson M, Almérás N, Dufresne SS, Robitaille J, Rhéaume C, Bujold E, et al. A 12-week exercise program for pregnant women with obesity to improve physical activity levels: an open randomised preliminary study. *PLoS One*. 2015;10(9):e0137742.
- 53 Goldstein RF, Abell SK, Ranasinha S, Misso M, Boyle JA, Black MH, et al. Association of gestational weight gain with maternal and infant outcomes: a systematic review and meta-analysis. *JAMA*. 2017;317(21):2207–25.
- 54 Marchi J, Berg M, Dencker A, Olander EK, Begley C. Risks associated with obesity in pregnancy, for the mother and baby: a systematic review of reviews. *Obes Rev*. 2015; 16(8):621–38.
- 55 American College of Obstetricians and Gynecologists’ Committee on Practice Bulletins—Obstetrics; Catalano PM, Koutrouvelis GO. collab. Practice bulletin: obesity in pregnancy. *Obstet Gynecol*. 2015;126(5): 1118–9.
- 56 Chu SY, Callaghan WM, Kim SY, Schmid CH, Lau J, England LJ, et al. Maternal obesity and risk of gestational diabetes mellitus. *Diabetes Care*. 2007;30(8): 2070–6.
- 57 Artal R. Exercise in pregnancy: guidelines. *Clin Obstet Gynecol*. 2016;59(3):639–44.
- 58 Tobias DK, Zhang C, Van Dam RM, Bowers K, Hu FB. Physical activity before and during pregnancy and risk of gestational diabetes mellitus: a meta-analysis. *Diabetes Care*. 2011;34(1):223–9.
- 59 Davenport MH, Nagpal TS, Mottola MF, Skow RJ, Riske L, Poitras VJ, et al. Prenatal exercise (including but not limited to pelvic floor muscle training) and urinary incontinence during and following pregnancy: a systematic review and meta-analysis. *Br J Sports Med*. 2018;52(21): 1397–404.
- 60 Magro-Malosso ER, Saccone G, Di Tommaso M, Roman A, Berghella V. Exercise during pregnancy and risk of gestational hypertensive disorders: a systematic review and meta-analysis. *Acta Obstet Gynecol Scand*. 2017; 96(8):921–31.
- 61 Ko YL, Chen CP, Lin PC. Physical activities during pregnancy and type of delivery in nulliparae. *Eur J Sport Sci*. 2016;16(3): 374–80.
- 62 Barakat R, Pelaez M, Lopez C, Montejó R, Coterón J. Exercise during pregnancy reduces the rate of cesarean and instrumental deliveries: results of a randomized controlled trial. *J Matern Neonatal Med*. 2012; 25(11):2372–6.
- 63 Santos PC, Maciel LYS, Abreu S, Mesquita AR, Mesquita CC, Lopes S, et al. Cultural adaptation and validation of the “pregnancy physical activity questionnaire” for the Portuguese population. *PLoS One*. 2023;18(1): e0279124.