

AUTOTRANSPLANTE RENAL BILATERAL COMO SOLUÇÃO PARA ANEURISMAS MÚLTIPLOS DOS RAMOS DAS ARTÉRIAS RENAI: RELATO DE CASO

BILATERAL KIDNEY AUTOTRANSPLANTATION AS A SOLUTION FOR MULTIPLE ANEURYSMS OF THE RENAL ARTERIES BRANCHES: A CASE REPORT

Daniel Mendes^{1*}, Rui Machado^{1,2}, Carlos Veiga¹, Carlos Veterano¹, Henrique Rocha¹, João Castro¹, Andreia Pinelo¹, Henrique Almeida¹, Rui de Almeida^{1,2}

1. Serviço de Angiologia e Cirurgia Vascular; Centro Hospitalar Universitário do Porto; Porto; Portugal
2. Instituto de Ciências Biomédicas Abel Salazar – Universidade do Porto; Porto; Portugal

Recebido em: 26/07/2021

Aceite para publicação em: 26/12/2022

RESUMO

Introdução: Os aneurismas da artéria renal são entidades clínicas raras cujos critérios de tratamento não se encontram totalmente estabelecidos.

O tratamento endovascular tem ganho aceitação no caso dos aneurismas do tronco da artéria renal. No entanto, no caso dos aneurismas dos ramos das artérias renais tais procedimentos não são possíveis devido à elevada morbilidade, pelo que não existe consenso quanto ao melhor tratamento. Apresentamos um caso clínico de uma doente com aneurismas múltiplos dos ramos das artérias renais, bilateralmente, tratados com reconstrução ex-vivo e autotransplante renal.

Apresentação do caso: Doente do sexo feminino de 35 anos diagnosticada com aneurismas das artérias renais bilateralmente durante a investigação de hipertensão arterial secundária. A doente foi submetida a tratamento cirúrgico sequencial com reconstrução ex-vivo e autotransplante renal tendo sido inicialmente tratado o rim direito. O procedimento cirúrgico iniciou-se pela nefrectomia laparoscópica transperitoneal. Procedeu-se à correção ex-vivo dos aneurismas “em banca” com aneurismectomia e aneurismorrafia. O enxerto foi implantado na fossa ilíaca direita com anastomoses às artéria e veia ilíacas externas sem intercorrências.

Três meses depois, foi realizado um procedimento idêntico à esquerda. À inspeção “em banca” observou-se um volumoso aneurisma na bifurcação da artéria renal e outro aneurisma de pequenas dimensões num dos ramos, ambos corrigidos com aneurismectomia e aneurismorrafia. O controlo angiográfico ex-vivo assegurou a preservação da patência das artérias do enxerto sem evidência de aneurismas. O rim foi implantado nos vasos ilíacos externos esquerdos sem intercorrências.

Foi realizado estudo com cintigrafia renal que confirmou a adequada função de ambos os rins. Aos nove meses de seguimento a doente apresenta-se assintomática com valores de creatinina sérica e taxa de filtração glomerular dentro dos valores normais.

Conclusão: O autotransplante renal com reparação arterial ex-vivo parece ser uma boa solução no tratamento de patologia aneurismática da artéria renal, nomeadamente, nos casos complexos com múltiplos aneurismas.

Palavras-chave

Reparo ex vivo da artéria renal; hipertensão; aneurisma da artéria renal; transplante de rim; autotransplante renal

*Autor para correspondência.

Correio eletrónico: daniel5.mds@gmail.com (D. Mendes).

ABSTRACT

Introduction: Renal artery aneurysms are rare clinical entities whose treatment criteria are not fully established. Endovascular treatment has gained acceptance in the case of renal artery trunk aneurysms. However, in the case of aneurysms of the renal artery branches, such procedures are often impossible due to high morbidity, so the question remains about the best treatment modality. We present a clinical case of multiple bilateral aneurysms of the renal artery branches in a young patient adequately treated with ex-vivo reconstruction and renal autotransplantation.

Case-presentation: A 35-year-old woman was diagnosed with bilateral renal artery aneurysms during an investigation of secondary arterial hypertension (AHT). The patient underwent surgical treatment with renal autotransplantation, and the right kidney was initially treated. Transperitoneal laparoscopic nephrectomy was performed. She underwent ex vivo repair of aneurysms, on the bench, with aneurysmectomy and aneurysmorrhaphy. The graft was implanted in the right iliac fossa with anastomoses to the external iliac artery and vein.

After three months, an identical procedure was performed on the left. Upon inspection, a large aneurysm was observed in the renal artery bifurcation and another small aneurysm in one of the branches; both corrected with aneurysmectomy and aneurysmorrhaphy. Ex-vivo angiographic control ensured the preservation of graft arteries patency without evidence of residual aneurysms. The kidney was implanted in the left external iliac vessels uneventfully.

A study with renal scintigraphy confirmed the adequate function of both kidneys. The patient is asymptomatic at nine months of follow-up with serum creatinine and glomerular filtration rate within normal values.

Conclusion: Renal autotransplantation with ex vivo arterial repair seems to be a good solution in treating aneurysms of the renal artery, particularly in complex cases with multiple aneurysms.

Keywords

Ex vivo renal artery repair; Hypertension; Renal artery aneurysm; Kidney transplantation; Renal autotransplantation

INTRODUCTION

Renal artery aneurysms are infrequent, occurring in about 0.1% of the population, increasing to 0.3% to 2.5% in angiographic and computed tomography studies.^(1,2) They are usually accidental findings on diagnostic imaging or are diagnosed in the study of secondary arterial hypertension. They can have different etiologies but are typically associated with fibromuscular dysplasia.⁽³⁾ The fact that fibromuscular dysplasia occurs more frequently on right-sided kidneys than on the left side could justify these aneurysms being commoner on the right side.⁽⁴⁾ Other etiologies include Behcet's disease, Von Recklinghausen's disease, arterial dissection, iatrogenic injury, and atherosclerosis, the latter in elderly patients.⁽⁵⁾ Usually asymptomatic, renal artery aneurysms may present with lower back or flank pain and hematuria. Also, it can present with renovascular hypertension mainly due to concomitant renal artery stenosis.⁽⁶⁾ Computed tomography angiography (CTA) is usually the diagnostic method used for suspected renal artery aneurysms. However, the visualization of aneurysms in the renal artery and hilar branches may be limited. In such cases, multi-angle catheter angiography may

be beneficial in facilitating the definition of the therapeutic strategy.⁽⁷⁾

There is still some controversy regarding the treatment criteria considering the growing evidence that this disorder has a benign behavior. There is some consensus that endovascular treatment may be a good option for aneurysms located in the trunk of the renal artery⁽⁸⁾. However, the evidence is not clear in cases of renal branch artery and multiple aneurysms. In these situations, ex-vivo reconstruction with cooling of the kidney offers the possibility of performing complex vascular reconstructions while preserving the functioning renal mass.

We present a complex case of bilateral renal artery branches multiple aneurysms. A detailed review of the clinical data was performed. Patients' informed consent was obtained for publication of the case report.

CASE-PRESENTATION

A 35-year-old woman with a history of arterial hypertension (AHT) being treated with three different classes of drugs has been exhaustively investigated in the suspicion of secondary AHT. No laboratory

abnormalities were found. The patient performed an abdominopelvic magnetic resonance angiography (MRA) to exclude renovascular hypertension, namely renal artery stenosis associated with fibromuscular dysplasia. This imaging exam revealed the presence of bilateral saccular renal artery aneurysms, measuring 24mm on the right kidney and approximately 19mm on the left kidney (FIGURE 1), with no evidence of morphologically significant stenosis in the renal arteries. No aneurysmal dilatations were observed in other arteries of the abdomen or pelvis.




Figure 1 Abdominal Magnetic Resonance Angiography (MRA) showing a right renal artery saccular aneurysm (arrow) measuring 24mm and a left renal artery saccular aneurysm (arrowhead) of approximately 19mm in diameter with no evidence of rupture or renal artery stenosis.

Bearing in mind that the aneurysms were located far from the ostium, apparently in a bifurcation site closer to the renal hilum, and to exclude multiple aneurysms, since bilateral aneurysms were observed, it was decided to perform a study by selective arteriography. Bilateral renal angiography confirmed the presence of saccular aneurysms of both renal arteries (FIGURE 2). Two saccular aneurysms were observed on the right, the largest with more than 20mm in diameter, and on the left, a saccular aneurysm with approximately 20mm. Similarly, no significant stenosis of the renal artery or its branches was observed.



Figure 2 Angiography revealing bilateral saccular renal artery aneurysms.

The patient underwent surgical treatment with ex vivo repair and total kidney autotransplantation at different times for both kidneys, about three months apart. The right kidney was initially treated. Transperitoneal laparoscopic nephrectomy was performed. A brief description of the technique includes the medial mobilization of the colon to access the retroperitoneum with the opening of the renal capsule. The dissection and isolation of the ureter, artery, and renal vein is carried out. The ureter is ligated with a clip as distally as possible, and the renal vessels are ligated using two Hem-o-lok clips. The kidney is extracted into an endobag through a small Gibson incision. After rapid extraction, the kidney is washed and perfused with Celsior® preservation solution. Warm ischemia time was of 2 minutes and 25 seconds. Upon visualization of the kidney in a bench, a large aneurysm in the renal bifurcation and another small saccular aneurysm in a first-order branch were observed. She underwent ex vivo repair of aneurysms with aneurysmorrhaphy, and angiographic control was performed (figure 3). For this purpose, the kidney was placed on a radiolucent table under the fluoroscopic arch. The renal artery was cannulated, and approximately 10cc of a mixture of equal parts of Visipaque®³²⁰ (Iodixanol) isosmolar, radiographic contrast medium, and Celsior® preservation solution was injected. The on-table angiography revealed the presence of a third aneurysm in another branch, corrected with aneurysmorrhaphy. Completion angiography confirmed the permeability of all branches and adequate correction of the aneurysms. The graft was implanted in the right iliac fossa in a heterotopic fashion through the same kidney extraction incision. After dissection and isolation of retroperitoneal iliac vessels, the renal vein was anastomosed to the external iliac vein in an end-to-side fashion. It was necessary to perform an elongation venoplasty using a spiral saphenous vein graft. The renal artery was connected to the external iliac artery in an end-to-side fashion with an interposition great saphenous vein graft. Creation of neoureterocystotomy according to Paquin's direct implantation technique was performed with insertion of a double J catheter. After seven days, the patient was discharged with normal renal function confirmed by renal scintigraphy. The double J catheter was removed after four weeks.

After three months, an identical procedure on the left side was performed with warm kidney ischemia of 2 minutes and 59 seconds. Upon inspection, a large



Figure 3 On the bench angiography after ex vivo repair of renal artery aneurysms before implantation on iliac vessels.

aneurysm was observed in the renal artery bifurcation and another small aneurysm in one of the branches; both corrected with aneurysmectomy and aneurysmorrhaphy. Ex-vivo angiographic control ensured the preservation of graft arteries patency without evidence of residual aneurysms. The kidney was uneventfully implanted in the left external iliac vessels (FIGURE 4).




Figure 4 Postoperative Computed Tomography Angiography (CTA) reveals properly implanted kidneys in both iliac fossae. A small area of infarction is noted at the upper pole of the left kidney. A double-J catheter in the ureter of the left kidney is observed considering the short time from surgical repair.

A study with renal scintigraphy confirmed the adequate function of both kidneys (FIGURE 5). The patient was discharged after seven days without complications. At nine months of follow-up, the patient is asymptomatic with serum creatinine values of 76.04 $\mu\text{mol/L}$ (IU), corresponding to a glomerular filtration rate of 86.1 mL/min/1.73 m². Pathological examinations of the aneurysms revealed a myxoid wall degeneration, with no other specific findings. Arterial hypertension resolved utterly, and it was possible to discontinue all antihypertensive drugs with blood pressure values within the reference ranges.




Figure 5 Renal scintigraphy, revealing kidneys located in the iliac fossa bilaterally with good perfusion and concentration of the radiopharmaceutical. Slight pyelocaliceal stasis is visualized, with slow radiopharmaceutical drainage.

DISCUSSION

Recent studies have revealed that renal artery aneurysms usually have a more benign clinical course than previously thought, with a slow growth rate. Although there are no sufficiently robust studies, most recommendations have been increasingly broad about dimensions for treatment.^(9, 10) Reported growth rates range from 0.65mm per year to 3mm per year, but the series with the highest number of patients points to less than 1mm per year.^(5, 11, 12)

Accepted indications for treatment include aneurysms larger than 30mm in diameter or symptomatic. Pregnancy has been associated with increased rupture rates, presumably secondary to hemodynamic changes and increased vascular volume and flow during pregnancy. Thus, the treatment of renal artery aneurysms in women of childbearing age, regardless of size, is justified, which was the case of our patient, who intended to become pregnant in the short term. Clinically relevant renal artery stenosis is present in 7% to 66% of renal artery aneurysm patients in all series. Although renal artery occlusive disease is not evidenced in all renal artery aneurysm patients with hypertension, it remains a valid indication for intervention.⁽⁹⁾ Our patient was one of these cases who had resistant hypertension without any evidence of stenosis or flow obstruction in the MRA and multi-angle selective angiography. The mechanism for hypertension in these patients is not fully established. However, it is thought that flow turbulence in the aneurysmal sac or eventual vascular compression causing dynamic stenosis may contribute. Aneurysmal sac thrombus embolization with cortical ischemia is another possible mechanism.^(5, 13)

Arteriography, previously considered the gold standard, was replaced by less invasive imaging tests in diagnosing and characterizing this pathology, such

as CTA and MRA. These exams allow an adequate assessment of the dimensions of the aneurysms, their anatomical location, and their relationship with adjacent structures. However, selective arteriography may be essential when there is a suspicion of more complex aneurysms, namely in distal or hilar aneurysms with involvement of the renal artery branches that may be lost on cross-sectional imaging.⁽²⁾ Our case demonstrated this, in which aneurysms of the renal artery branches were only visualized on arteriography, which was essential in therapeutic planning.

Several therapeutic options are available, however, choosing the most appropriate treatment remains controversial. With the increasing use of endovascular techniques, simple and proximal renal artery trunk aneurysms could be treated by placing a stent-graft, selective coil embolization or balloon-assisted coil embolization with reported success rates of 73% to 100% and a morbidity rate of 13% to 60%.^(2,14,15) Postembolization syndrome, defined as a combination of fever, leukocytosis, abdominal pain, nausea, and vomiting, was the most common reported complication.⁽¹⁶⁾ Endovascular intervention in more complex renal artery aneurysms results in high incidences of the post-embolization syndrome and segmental renal ischemia. Also, there are uncertain long-term data.⁽⁵⁾ In more complex aneurysms, surgical repair is the treatment of choice. Several techniques are described and include in-situ repair, ex-vivo reconstruction with partial (without division of the ureter) or total kidney autotransplantation, and primary nephrectomy.

In situ repair can be performed by conventional open, laparoscopic, or robotic approach.⁽¹⁷⁾ Usually used for single aneurysm repair, in situ repair is not recommended in complex aneurysms of bifurcations or branches or multiple aneurysms when a long ischemia time is expected to perform the vascular reconstruction.⁽¹⁸⁾ Available conventional in situ reconstructions include aneurysm resection with primary angioplasty (with or without branch reimplantation), patch angioplasty, primary anastomosis, interposition bypass, aortorenal bypass, splanchnic-renal bypass, and small aneurysm plication.

Although historically treated with nephrectomy, current data support that complex distal branches, or multiple aneurysms are best addressed with ex-vivo repair and renal autotransplantation, providing durable repair and maximum graft protection. This technique somehow increases with less invasive procedures for kidney harvest. Laparoscopic living donor nephrectomy reduced hospital stay, pain, and time to return to full function.⁽¹⁹⁾ However, availability and experience should be the decisive factors. At our center, experienced laparoscopic urologists are

integrated into the surgical teams to perform autotransplant procedures, so we use the laparoscopic technique whenever possible.

Although direct studies comparing in situ reconstructions and ex-vivo repairs have not been performed, better patency and renal viability rates have been observed in ex-vivo reconstructions.^(8,20)

Murray et al. describe the most extensive series with a 92% success rate with 12 patients treated without significant mortality or morbidity.⁽²¹⁾ Gallagher et al. reported seven ex-vivo reconstructions after laparoscopic nephrectomy for complex aneurysmal disease to avoid surgical incision morbidity; these authors described excellent technical success, no mortality, no ureteral morbidity, and a 28% incidence of perioperative morbidity.⁽²²⁾ Duprey et al. showed primary patency at 30 days of 90.8% with a survival of 95% at nine years.⁽⁵⁾ Data from our center with the kidney autotransplantation technique were recently published with a kidney patency rate of 93% with a follow-up of 47.2 months and no mortality.⁽¹²⁾ As in this case, we use laparoscopic surgery to harvest the kidney in most cases due to the excellent expertise of the surgical team, decreasing surgical aggressiveness and morbidity. Primary nephrectomy is currently a treatment only used in extreme cases of multiple complex aneurysms, refractory hypertension, or when a reconstructive surgical solution cannot be performed, namely after the failure of endovascular treatment.⁽¹⁸⁾ In case of a ruptured renal artery aneurysm with hemodynamic instability, it may be an option. However, all efforts must be made to preserve the functioning renal parenchyma.

CONCLUSION

In our experience, renal autotransplantation with ex vivo arterial repair seems to be a good solution in treating multiple and complex renal artery aneurysms, with good mid and long-term results. The results in the literature corroborate this evidence with a high success rate, low morbidity and mortality, high patency rates and preservation of renal function.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

AUTOTRANSPLANTE RENAL BILATERAL COMO SOLUÇÃO PARA ANEURISMAS MÚLTIPLOS DOS RAMOS DAS ARTERIAS RENAIS: RELATO DE CASO

1. Stanley JC, Rhodes EL, Gewertz BL, Chang CY, Walter JF, Fry WJ. Proceedings: Renal artery aneurysms: significance of macroaneurysms exclusive of dissections and fibrodysplastic mural dilatations. *J Cardiovasc Surg (Torino)* 1976; 17(1): 85.
2. Coleman DM, Stanley JC. Renal artery aneurysms. *J Vasc Surg* 2015; 62(3): 779-85.
3. Laser A, Flinn WR, Benjamin ME. Ex vivo repair of renal artery aneurysms. *J Vasc Surg* 2015; 62(3): 606-9.
4. Augustin G, Kulis T, Kello N, Ivkovic V. Ruptured renal artery aneurysm in pregnancy and puerperium: literature review of 53 cases. *Arch Gynecol Obstet* 2019; 299(4): 923-31.
5. Duprey A, Chavent B, Meyer-Bisch V, et al. Editor's Choice - Ex vivo Renal Artery Repair with Kidney Autotransplantation for Renal Artery Branch Aneurysms: Long-term Results of Sixty-seven Procedures. *Eur J Vasc Endovasc Surg* 2016; 51(6): 872-9.
6. Tham G, Ekelund L, Herrlin K, Lindstedt EL, Olin T, Bergentz SE. Renal artery aneurysms. Natural history and prognosis. *Ann Surg* 1983; 197(3): 348-52.
7. Liu PS, Platt JF. CT angiography of the renal circulation. *Radiol Clin North Am* 2010; 48(2): 347-65, viii-ix.
8. Henke PK, Cardneau JD, Welling TH, 3rd, et al. Renal artery aneurysms: a 35-year clinical experience with 252 aneurysms in 168 patients. *Ann Surg* 2001; 234(4): 454-62; discussion 62-3.
9. Chaer RA, Abularrage CJ, Coleman DM, et al. The Society for Vascular Surgery clinical practice guidelines on the management of visceral aneurysms. *J Vasc Surg* 2020; 72(1S): 3S-39S.
10. Tsilimparis N, Reeves JG, Dayama A, Perez SD, Debus ES, Ricotta JJ, 2nd. Endovascular vs open repair of renal artery aneurysms: outcomes of repair and long-term renal function. *J Am Coll Surg* 2013; 217(2): 263-9.
11. Klausner JQ, Lawrence PF, Harlander-Locke MP, et al. The contemporary management of renal artery aneurysms. *J Vasc Surg* 2015; 61(4): 978-84.
12. Machado M, Machado R, Almeida R. Renal Autotransplantation for The Treatment of Renal Artery Aneurysm. *Ann Vasc Surg* 2021.
13. Lumsden AB, Salam TA, Walton KG. Renal artery aneurysm: a report of 28 cases. *Cardiovasc Surg* 1996; 4(2): 185-9.
14. Elaassar O, Auriol J, Marquez R, Tall P, Rousseau H, Joffre F. Endovascular techniques for the treatment of renal artery aneurysms. *Cardiovasc Intervent Radiol* 2011; 34(5): 926-35.
15. Sedat J, Chau Y, Baque J. Endovascular treatment of renal aneurysms: a series of 18 cases. *Eur J Radiol* 2012; 81(12): 3973-8.
16. Zhang Z, Yang M, Song L, Tong X, Zou Y. Endovascular treatment of renal artery aneurysms and renal arteriovenous fistulas. *J Vasc Surg* 2013; 57(3): 765-70.
17. Giulianotti PC, Bianco FM, Addeo P, Lombardi A, Coratti A, Sbrana F. Robot-assisted laparoscopic repair of renal artery aneurysms. *J Vasc Surg* 2010; 51(4): 842-9.
18. Contarini E, Takagi K, Kimenai H, et al. Kidney Autotransplantation for Renal Artery Aneurysm: Case Series and a Systematic Review. *Ann Vasc Surg* 2021; 77: 349 e5- e18.
19. Nanidis TG, Antcliffe D, Kokkinos C, et al. Laparoscopic versus open live donor nephrectomy in renal transplantation: a meta-analysis. *Ann Surg* 2008; 247(1): 58-70.
20. Pfeiffer T, Reiher L, Grabitz K, et al. Reconstruction for renal artery aneurysm: operative techniques and long-term results. *J Vasc Surg* 2003; 37(2): 293-300.
21. Murray SP, Kent C, Salvatierra O, Stoney RJ. Complex branch renovascular disease: management options and late results. *J Vasc Surg* 1994; 20(3): 338-45; discussion 46.
22. Gallagher KA, Phelan MW, Stern T, Bartlett ST. Repair of complex renal artery aneurysms by laparoscopic nephrectomy with ex vivo repair and autotransplantation. *J Vasc Surg* 2008; 48(6): 1408-13.