

Radiological Case Report / Caso Clínico

Imaging Features of Bothropic Accident

Achados de Imagem de Acidente Botrópico

Artur Ramos Sarmet dos Santos¹, Rafaela Queiroz de Morais¹, Cristina Asvolinsque Pantaleão Fontes¹, Diogo Goulart Corrêa¹

¹Department of Radiology, Federal Fluminense University, Niterói, Brazil

Address

Artur Ramos Sarmet dos Santos
Department of Radiology
Federal Fluminense University
Avenida Marquês de Paraná, 303, Centro
Niterói, RJ, Brazil. Zip Code: 24070-035
e-mail: a.sarmet@hotmail.com

Received: 27/08/2021

Accepted: 22/11/2021

Published: 13/01/2022

© Author(s) (or their employer(s)) and ARP
2021. Re-use permitted under CC BY-NC. No
commercial re-use.

Abstract

Snakebite envenoming is a neglected public health issue, which is potentially life-threatening, caused by toxins present in the bite of a venomous snake. Bothropic venom has important physiopathological effects, with local lesions and proteolytic action, as well as coagulation cascade activation, consumption of fibrinogen, leading to hemorrhagic events. In this article, we present the case of a 61-year-old man who was bitten by a snake on his left thigh, and evolved with hypotension, lethargy, nausea, and vomiting. Computed tomography of the brain, chest, abdomen, and pelvis demonstrated intracerebral hemorrhage in the left frontal lobe, bilateral subarachnoid hemorrhage, moderate left hemothorax and soft tissue edema at the site of the snakebites on the left thigh. Despite adequate treatment, the patient died on the fourth day of hospitalization. It is important that the emergency radiologist correctly recognize the imaging findings and the ophidian accident as the cause of the hemorrhagic manifestations.

Keywords

Bothropic accident; Computed tomography; Intracranial hemorrhage; Hemothorax.

Resumo

A mordedura de cobra é uma questão de saúde pública negligenciada, potencialmente fatal, causada por toxinas presentes no venoso da cobra. O veneno botrópico tem importantes efeitos fisiopatológicos, com lesões locais e ação proteolítica, bem como ativação da cascata da coagulação, consumo de fibrinogênio, levando a eventos hemorrágicos. Neste artigo, apresentamos o caso de um homem de 61 anos que foi mordido por uma cobra na coxa esquerda, e evoluiu com hipotensão, letargia, náuseas e vômitos. A tomografia computadorizada do crânio, tórax, abdome e pelve demonstrou hemorragia intracerebral no lobo frontal esquerdo, hemorragia subaracnóidea bilateral, hemotórax esquerdo moderado e edema nas partes moles, no local das mordeduras da cobra na coxa esquerda. Apesar do tratamento adequado, o paciente morreu no quarto dia de hospitalização. É importante que o radiologista da emergência reconheça corretamente os achados das imagens e o acidente ofídico como causa das manifestações hemorrágicas.

Palavras-chave

Acidente botrópico; Tomografia computadorizada; Hemorragia intracraniana; Hemotórax.

Introduction

Snakebite envenoming is a potentially life-threatening disease caused by toxins in the bite of a venomous snake. Furthermore, it is a neglected public health issue and a common medical emergency in many tropical countries.¹ According to the World Health Organization, about 5.4 million snakebites occur each year, resulting in 1.8 to 2.7 million cases of envenomings (poisoning from snake bites). There are between 81,410 and 137,880 deaths and around three times as many amputations and other permanent disabilities each year secondary to these events.² Tropical areas have the highest prevalence of venomous snakes, including Southeast Asia, Southwestern Europe, Sub-Saharan Africa, the United States of America and Latin America (Figure 1).³

The Viperidae family is responsible for most of the registered snakebite accidents in Latin America. In Brazil, the genus *Bothrops* is responsible for 85% of the ophidian envenomation.⁴ Bothropic venom has important physiopathological effects, with local lesions and tissue necrosis (proteolytic action). It also activates the coagulation cascade and promotes blood incoagulability through the

consumption of fibrinogen, leading to hemorrhagic events by causing damage to the basal membrane of capillaries associated with thrombocytopenia (anticoagulant action). Moreover, bothropic venom promotes the release of hypotensive substances that, associated with the hemorrhagic manifestations, configure a life-threatening situation, which are frequent in this type of ophidian accident.⁵




Figure 1 – Global distribution of venomous snakes. Adapted from: <https://apps.who.int/bloodproducts/snakeantivenoms/database>.

Case Presentation

A 61-year-old man was bitten by a snake on his left thigh in Macaé, a city in the countryside of Rio de Janeiro, Brazil, and was taken to the emergency department of a tertiary hospital. The patient arrived with an estimated time of 5 hours after the ophidic accident and presented with hypotension, lethargy, with sphincter release, nausea and vomiting. Besides the edema at the site of the snakebite, several snakebite marks were noted in the left thigh. The number of bites raised the alert for the possibility of a large amount of venom inoculated, being a great predictor of severity.

The snake was identified as the *Bothrops atrox* (common lancehead) by the patient's relatives, who reported finding him unconscious, justifying the number of bites. A few minutes after admission to the hospital, he presented a generalized tonic-clonic seizure, with sphincter release, melena, hemoptysis and tachypnea, requiring orotracheal intubation and ventilatory support. On physical examination, the patient presented signs of circulatory shock, in addition to lethargy and abolished breath sound on chest auscultation in the left lung base. Computed tomography (CT) of the brain, chest, abdomen and pelvis were performed, and demonstrated intracerebral hemorrhage in the left frontal lobe, bilateral subarachnoid hemorrhage, moderate left hemothorax and soft tissue edema at the site of the snakebites on the left thigh (Figure 2).

threatening manifestations are hypotension, circulatory shock, acute kidney injury, and hemorrhagic events. Complications, although infrequent, are almost always fatal and difficult to manage and include abscess formation, necrosis, and compartment syndrome.⁶

The main effects of the *Bothrops* venom are related with coagulation disorders, hemorrhage and local tissue damage. Most of bothropic venoms are composed by metalloproteases, which are responsible for pro- and anti-coagulant disorders and hemorrhage; Serine proteases, which affect different factors involved in the coagulation cascade; Phospholipases A2, which have myotoxic, and neurotoxic effects, generate edema, platelet aggregation disturbance, and tissue damage, among others effects; L-amino acid oxidases, that induce the release of H₂O₂, with cytotoxic effects, as well as induction of hemorrhage and apoptosis, and inhibition of platelet aggregation; and C-type lectins, which disrupts blood homeostasis by inducing/inhibiting platelet aggregation or activating/consuming coagulation factors.⁷ *Bothrops* present an ontogenetic shift in its venom composition. The venom of adult specimens shows predominantly proteolytic activity and local tissue damage, whereas the venom of juveniles causes mainly hemorrhagic disorders.^{4,6} Although in the case presented in this study, we did not recover the snake responsible for the accident, the great hemorrhagic manifestations make us believe that the accident was caused by a young snake.

Although ophidic accidents have a certain frequency worldwide, the most common is the presence of a single



Figure 2 – (A) Brain CT-scan without contrast showing intraparenchymal hemorrhage with perilesional edema in the left frontal lobe (red arrow) and bilateral subarachnoid hemorrhage (yellow arrows). (B) Axial nonenhanced chest CT (mediastinal window) presenting moderate left pleural effusion with high density (50 UH) compatible with hemothorax (red arrow). (C) Pelvic CT-scan without contrast demonstrating soft tissue edema with enlargement of the quadriceps muscles (red arrow) and densification of the subcutaneous adipose tissue (green arrow) at the site of the snakebite on the left thigh.

His blood tests presented low platelet count (20.000 cells/mm³), leukocytosis (17.480 cells/mm³), high creatine kinase (CK: 870 units per litre), elevated levels of international normalized ratio (INR: 3,5) and activated partial thromboplastin (aPTT: 80 seconds). Despite treatment with botropic antivenom being promptly instituted, along with all necessary clinical support measures, the patient suffered a cardiac arrest refractory to cardiopulmonary resuscitation measures, on the fourth day of hospitalization.

Discussion

This is a case of a patient developing severe coagulopathy disorder and clinical manifestations of cerebral hemorrhage and acute hypoxaemic respiratory failure after being bitten a few times by *Bothrops atrox*. Local manifestations of bothropic accidents include pain, edema, often associated with ecchymosis and local bleeding. Lymphadenopathy and blisters may appear in the process.^{3,5} Systemic and life-

snakebite mark, with two being very uncommon and several being extremely rare, drawing our attention to this case and serving as a predictor of severity. In addition, the hemorrhagic manifestations of this case have important differential diagnoses often seen in emergency departments, such as intracranial hemorrhage secondary to ruptured aneurysms, and hemothorax from chest trauma.

Conclusion

This case provides imaging findings including intraparenchymal hemorrhage, subarachnoid hemorrhage, hemothorax, and signs of soft tissue edema at the site of the snakebite, all at once. Therefore, it is essential for the emergency physician to be aware of imaging features of the hemorrhagic manifestations of ophidic accidents, to take the correct therapeutic approach, such as neuroprotective measures.

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Supporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Agarwal R, Aggarwal AN, Gupta D. Elapid snakebite as a cause of severe hypertension. *J Emerg Med.* 2006;30:19-320.
2. da Silva Souza A, de Almeida Gonçalves Sachett J, Alcântara JA, et al. Snakebites as cause of deaths in the Western Brazilian Amazon: why and who dies? Deaths from snakebites in the Amazon. *Toxicon.* 2018;145:15-24.
3. Pinho FM, Burdmann EA. Fatal cerebral hemorrhage and acute renal failure after young Bothrops jararacussu snake bite. *Ren Fail.* 2001;23:269-77.
4. Hatakeyama DM, Tasima LJ, Bravo-Tobar CA, et al. Venom complexity of bothrops atrox (common lancehead) siblings. *J Venom Anim Toxins Incl Trop Dis.* 2020;26:e20200018.
5. Tinoco HB, Norberg AN, Pile E, Carvalho CRP, Silva DA, Guerra-Sanches F. Snake envenomations in northwest counties of the Rio de Janeiro State, Brazil. *J Venom Anim Toxins incl Trop Dis.* 2005;11:34-8.
6. Longbottom J, Shearer FM, Devine M, et al. Vulnerability to snakebite envenoming: a global mapping of hotspots. *Lancet.* 2018;392:673-84.
7. Hatakeyama DM, Jorge Tasima L, da Costa Galizio N, et al. From birth to adulthood: an analysis of the Brazilian lancehead (bothrops moojeni) venom at different life stages. *PLoS One.* 2021;16:e0253050.