

## ORIGINAL ARTICLE

# Occupational Chromium Allergy and Contact Dermatitis: 10 Years Evolution

## *Dermatite de Contacto Alérgica Ocupacional ao Crómio: 10 Anos de Estudo*

Received/Recebido  
2021/02/16Accepted/Aceite  
2021/02/28Published/Publicado  
2021/09/30Alexandre Afonso<sup>1\*</sup>, Joana Silva<sup>1\*</sup>, Ricardo Silva<sup>1</sup>, Vânia Pacheco<sup>1</sup>, Vitor Pinheiro<sup>1</sup>, Isabel Antunes<sup>1</sup>, Margarida Gonçalves<sup>2</sup><sup>1</sup>Serviço de Saúde Ocupacional do Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal<sup>2</sup>Serviço de Dermatovenerologia do Centro Hospitalar e Universitário de Coimbra e Faculdade de Medicina da Universidade de Coimbra, Coimbra, Portugal

\* Joint first authors - equally contribution / Co-primeiros autores - contribuíram igualmente para o estudo

**ABSTRACT – Introduction:** Chromium has been responsible for occupational skin disease, both irritant and/or allergic contact dermatitis. In order to minimize this problem, regulatory measures limiting its use have been implemented. The impact of these regulations should be monitored over time. **Material and Methods:** A retrospective study from 2009 to 2018 was carried out in the Department of Dermatology to assess the temporal trend of chromium sensitization evaluated by patch testing with potassium dichromate 0.5% pet, particularly considering age, sex and its relationship with occupational activity.

**Results:** 3277 individuals were studied, 2369 females (72.29%) and 908 males (27.71%). Of those 621 (18.95%) fulfilled criteria for occupational contact dermatitis.

Chromium allergy was found in 118 (3.60%) patients, 64 females (54.4%) and 54 males (45.76%). We found that male patients were significantly more frequent (45.76% vs 27.03%;  $p < 0.001$ ), as well as occupational dermatitis (31.35% vs 18.47%;  $p < 0.001$ ), hand dermatitis (53.54% vs 23.52%;  $p < 0.001$ ) and age above 40 years (70.34% vs 56.76%;  $p = 0.003$ ).

**Discussion/Conclusion:** The results show a high prevalence (3.60%) of chromium positive patch tests. In our study, there has not been a significant decrease in sensitization to chromium over the years. Interestingly cases classically related to this allergy (cement in the building industry) are no longer the main cause of the problem. We found other sources such as leather as a main cause of allergic sensitization, suggesting a change of the panorama.

**KEYWORDS –** Chromium/adverse effects; Dermatitis, Allergic Contact; Dermatitis, Occupational; Patch Tests.

**RESUMO – Introdução:** O crómio tem sido identificado como responsável por dermatites de contato irritativas ou alérgicas ocupacionais. Para minimizar o seu impacto, foram implementadas medidas regulatórias que limitam o seu uso, cujo impacto deve ser monitorizado ao longo do tempo.

**Material e Métodos:** Estudo retrospectivo realizado entre 2009 e 2018 no Serviço de Dermatologia dos CHUC para avaliar a tendência temporal da sensibilização ao crómio em testes epicutâneos com dicromato de potássio a 0,5% vas, de acordo com a idade, sexo e sua relação ocupacional.

**Resultados:** Foram estudados 3277 indivíduos, 2369 mulheres (72,29%) e 908 homens (27,71%), dos quais 621 (18,95%) tinham critérios para dermatite de contato ocupacional.

A sensibilização ao crómio foi encontrada em 118 (3,60%) pacientes, 64 mulheres (54,4%) e 54 homens (45,76%). Destes, o sexo masculino foi significativamente mais frequente (45,76% vs 27,03%;  $p < 0,001$ ), assim como dermatite ocupacional (31,35% vs 18,47%;  $p < 0,001$ ), dermatite das mãos (53,54% vs 23,52%;  $p < 0,001$ ) e idade acima de 40 anos (70,34% vs 56,76%;  $p = 0,003$ ).

**Discussão / Conclusão:** Os resultados mostram uma alta prevalência (3,60%) de sensibilização ao crómio. Na nossa amostra, não houve diminuição estatisticamente significativa da sensibilização ao crómio ao longo dos anos, contudo, a elevada prevalência de outras fontes de exposição, como no couro, sugere que o foco é agora outro diferente dos casos classicamente relacionados a essa alergia (cimento na construção civil).

**PALAVRAS-CHAVE –** Crómio/efeitos adversos; Dermatite Alérgica de Contato; Dermatite Ocupacional; Testes Epicutâneos.

## INTRODUCTION

Occupational contact dermatitis (OCD), which encompasses irritant, allergic, and immediate reactions, is a limiting factor of professional activities and a cause of discomfort that can lead to decreased work capacity and absenteeism.<sup>1,2</sup> OCD represents the main occupational skin disease (up to 70% of cases)<sup>3-5</sup> and is also one of the main occupational diseases.

Chromium has been responsible for occupational skin disease, both irritant and/or allergic contact dermatitis. Industrialization and new materials used currently have resulted in increased exposure to chromium ions.<sup>6,7</sup> The main sources of exposure to chromium include cement, leather products, anti-corrosive paints, cleaning products, metal alloys, cosmetics, mobile phone components, implants / prosthetics, among others.<sup>1,7-12</sup>

The history of chromium as an allergen has several decades and

is associated with different interventions to minimize exposure to this metal, which have managed to gradually change the panorama of the epidemiology of chromium allergy.<sup>6,7,12-16</sup> In Denmark, in 1983, the implementation of obligation to add ferrous sulphate to cement reduced the concentration of water-soluble chromium to values <2 ppm. Then Finland in 1987 and Sweden in 1989 implemented the same measure<sup>7</sup> and, in 2003, the European directive restricting also marketing and use of cement containing hexavalent chromium in concentrations > 2 ppm came into force,<sup>17</sup> a directive that was taken into action in Portugal in 2005.<sup>18</sup>

In 2003 the concentration of hexavalent chromium in leather, namely in protective gloves, was limited to <10 ppm and in 2009 further reduced to <3 ppm. The decision to limit hexavalent chromium below 3ppm in leather products that contact the skin was finally adopted by the European Commission in 2015.<sup>7,17</sup>

It is interesting to note that contact allergy to chromium is usually concomitant with other allergies, especially other metals.<sup>12</sup> However this is not due to cross-reactivity but mostly to the synchronous presence of other metals in sensitizing products.<sup>7</sup> The aim of the study is to carry out an epidemiological analysis of the prevalence of contact allergy to chromium and evaluate if the regulatory measures implemented so far in Portugal have been enough to lessen the problem of sensitization to chromium.

## METHODS

This is a retrospective study conducted for 10 years (2009 to 2018) at the Cutaneous Allergology of the Department of Dermatology, Hospital and University Center of Coimbra (Portugal) and involved all consecutive patients who underwent epicutaneous patch tests for the study of suspected allergic contact dermatitis (ACD) or other delayed hypersensitivity reactions with skin involvement.

All patients who had a reactive epicutaneous test for chromium were included and the following parameters evaluated: gender, age, personal history of atopy, main location of the lesions, time of evolution of the lesions, occupation, other positive allergens (namely other metals and rubber allergens usually related to concomitant exposure in an occupational setting), clinical and occupational relevance. The population with chromium allergy was compared with the general population tested by the MOAHLFA index (male, occupational dermatitis, atopic dermatitis, hand dermatitis, leg ulcer/dermatitis, face dermatitis and age > 40 years).

### Epicutaneous tests

All patients were tested with the European baseline and additional series, according to the medical history and the tasks they developed. Allergens from Chemotechnique (Chemotechnique Diagnostics®, Vellinge, Sweden) or Trolab allergens® (Almirall GmbH, Germany) were applied for 48 hours on the back using Finn Chambers® (Epitest Ld, Almirall) or IQ Chambers® (Chemotechnique Diagnostics®, Vellinge, Sweden). Readings were performed at day 2-3 and day 4-7, according to European Society of Contact Dermatitis (ESCD) guidelines.<sup>5</sup> Epicutaneous tests or open tests were sometimes carried out with products brought by the patient, collected in the workplace or in his personal environment. Reactions were considered positive if at least erythema and infiltration were observed (1+ or more intense). Positive reactions were interpreted as having current, past or unknown relevance or explained by cross-reactions.

## Statistical analysis

For the statistical analysis, the program IBM SPSS Statistics for Windows, version 22.0 was used. Quantitative variables were tested for normality using Shapiro Wilk test. Correlation between variables were analyzed with Pearson's or Spearman's Correlation Coefficient. The X2 test of independence was used to compare categorical variables. Fisher's exact test was applied when the expected value in any of the cells of contingency table was < 5. Significance level was established at 0.05.

## RESULTS

Between 2009 and 2018, 3277 individuals were studied, 2369 females (72.29%) and 908 males (27.71%), 621 of whom (18.95%) with criteria for occupational contact dermatitis.

Chromium allergy was found in 118 (3.60%) patients, 64 females (54.4%) and 54 males (45.76%), representing respectively 2.70% and 5.95% of all the women and men tested in this period. The average age of females was 46.63±14.67 and males 54.06±13.13 years ( $p=0.004$ ).

No statistically significant correlation was found in the number of cases over time ( $\rho=-0,265$ ;  $p>0,05$ ). This remained true, when performing the analysis for age groups and gender (Table 1).

When comparing the MOAHLFA index of patients with and without chromium allergy, we found that among chromium sensitized patients the percentage of male patients was statistically more frequent (45.76% vs 27.03%;  $p<0.001$ ), as well as occupational dermatitis (31.35% vs 18.47%;  $p<0.001$ ), hand dermatitis (53.54% vs 23.52%;  $p<0.001$ ) and age above 40 years (70.34% vs 56.76%;  $p=0.003$ ).

The history of atopy was found in 31 patients with chromium allergy (asthma in 8, rhinitis in 4, atopic dermatitis in 13, asthma and rhinitis in 4, asthma and atopic dermatitis in 1 and rhinitis and atopic dermatitis in 1). This conveys for a similar proportion between those with and without chromium allergy (26.27% vs 30.83%,  $p > 0.05$ ). Likewise face dermatitis and leg ulcer/stasis dermatitis had no statistical difference between chromium positive and chromium negative patients (Table 2).

Table 3 shows the MOAHLFA index of all patients with chromium allergy stratified by gender. Occupational dermatitis was more frequent in men (40.74% vs 23.44%;  $p=0.044$ ) and males also had more frequent hand dermatitis (66.67% vs 40.63%;  $p=0.005$ ).

Regarding the distribution of cases by age and gender, the frequency was increased in the female gender in the 21-40 age group (37.50% vs 14.81%;  $p=0.006$ ), but superior in men older than 60 ( $p=0.006$ ). In the remaining age groups, we did not find significant differences (Table 4).

In this 10-year interval we also found 210 (6.41%) cases of cobalt allergy and 858 (26.18%) cases of nickel allergy. Ninety-eight (2.99%) patients were positive to thiuram mix, 70 (2.14%) to carba mix, 17 (0.52%) for mercapto mix and 20 (0.61%) for mercaptobenzothiazole.

In the subgroup of patients with chromium allergy, a concomitant positive patch test to cobalt occurred in 58 (49.15%), to nickel in 68 (57.63%), thiuram mix in 24 (20.34%), carba mix in 15 (12.71%), mercapto mix in 5 (4.24%) and mercaptobenzothiazole in 8 (6.78%). Simultaneous positive tests for chromium, cobalt and nickel were found in 41 (34.75%) patients.

In crude analysis, comparing chromium positive patients with

**Table 1** - Distribution of positive tests to potassium dichromate (Cr) by year and gender among all patchtested patients.

	Total tests n (M/F)	Total Cr + n (%)	Cr+ Men n (%)	Cr + Woman n (%)
2009	342 (96/246)	15 (4.39%)	4 (4.17%)	11 (4.47%)
2010	296 (71/225)	15 (5.07%)	9 (12.67%)	6 (2.67%)
2011	276 (82/196)	13 (4.71%)	4 (4.88%)	9 (4.59%)
2012	337 (92/245)	5 (1.48%)	3 (3.26%)	2 (0.82%)
2013	301 (73/228)	14 (4.65%)	3 (4.10%)	11 (4.82%)
2014	310 (86/224)	4 (1.29%)	3 (3.49%)	1 (0.45%)
2015	339 (99/240)	11 (3.24%)	8 (8.08%)	3 (1.25%)
2016	343 (103/240)	13 (3.80%)	8 (7.77%)	5 (2.08%)
2017	372 (110/262)	14 (3.76%)	5 (4.55%)	9 (3.43%)
2018	361 (96/265)	14 (3.88%)	7 (7.29%)	7 (2.64%)
<b>Total studied</b>	<b>3277 (908/2369)</b>	<b>118 (3.60%)</b>	<b>54 (5.95%)</b>	<b>64 (2.70%)</b>

**Table 2** - Comparison of the MOAHLFA index between patients with a positive test to chromium and chromium negative test in the tested sample.

	Total (%)	Chromium positive (%)	Chromium negative (%)	p-value*	Comparison
<b>Male</b>	908 (27.71%)	54 (45.76%)	854 (27.03%)	<0.001	↑
<b>Occupational</b>	621 (18.95%)	37 (31.35%)	584 (18.47%)	<0.001	↑
<b>Hand dermatitis</b>	805 (24.57%)	62 (53.54%)	743 (23.52%)	<0.001	↑
<b>Atopy</b>	1005 (30.67%)	31 (26.27%)	974 (30.83%)	0.291	↔
<b>Leg ulcer/stasis dermatitis</b>	205 (6.26%)	4 (3.39%)	201 (6.36%)	0.190	↔
<b>Face dermatitis</b>	839 (25.60%)	22 (18.64%)	817 (25.86%)	0.078	↔
<b>Age &gt; 40</b>	1876 (57.25%)	83 (70.34%)	1793 (56.76%)	0.003	↑
<b>Total studied</b>	<b>3277</b>	<b>118</b>	<b>3159</b>		

\* X<sup>2</sup> test comparing chromium positive cases with chromium negative cases.

**Table 3** - MOHALFA index of chromium positive patients stratified by gender.

	Total n (%)	Men n (%)	Women n (%)	p-value*	Comparison
<b>Male</b>	54 (45.76%)	-	-	-	-
<b>Occupational</b>	37 (31.35%)	22 (40.74%)	15 (23.44%)	0.044	↑
<b>Hand dermatitis</b>	62 (53.54%)	36 (66.67%)	26 (40.63%)	0.005	↑
<b>Atopy</b>	31 (26.27%)	11 (20.37%)	20 (31.25%)	0.181	↔
<b>Leg ulcer/stasis dermatitis</b>	4 (3.39%)	2 (3.70%)	2 (3.13%)	0.624	↔
<b>Face dermatitis</b>	22 (18.64%)	9 (16.67%)	13 (20.31%)	0.612	↔
<b>Age &gt; 40</b>	83 (70.34%)	44 (81.48%)	39 (60.94%)	0.015	↑
<b>Total studied</b>	<b>118</b>	<b>54</b>	<b>64</b>		

\* X<sup>2</sup> test comparing chromium positive cases between men and women.

**Table 4** - Distribution by age group and gender.

	Total (%)	Men (%)	Women (%)	p-value*
0-20	1 (0.85%)	0	1 (1.56%)	-
21-40	32 (27.12%)	8 (14.81%)	24 (37.50%)	0.006
41-60	58 (49.15%)	28 (51.85%)	30 (46.88%)	-
61+	27 (22.88%)	18 (33.33%)	9 (14.06%)	0.006
<b>Total studied</b>	<b>118</b>	<b>54</b>	<b>64</b>	

\* X<sup>2</sup> test comparing chromium positive cases by gender age group.

**Table 5** - Concomitant positive patch test reactions to other metals and rubber allergens in patients reacting to chromium versus non-reactive patients.

	Total n (%)	Chromium positive (%)	Chromium negative (%)	p-value*	Comparison
Cobalt	210 (6.41%)	58 (49.15%)	152 (4.81%)	<0.001	↑
Nickel	858 (26.18%)	68 (57.63%)	790 (25.01%)	<0.001	↑
Thiuram mix	98 (2.99%)	24 (20.34%)	74 (2.34%)	<0.001	↑
Carbamates mix	70 (2.14%)	15 (12.71%)	55 (1.74%)	<0.001	↑
Mercapto mix	17 (0.52%)	5 (4.24%)	12 (0.38%)	<0.001	↑
Mercapto benzothiazole	20 (0.61%)	8 (6.78%)	12 (0.38%)	<0.001	↑
<b>Total studied</b>	<b>3277</b>	<b>118</b>	<b>3159</b>		

\* X<sup>2</sup> test comparing chromium positive cases with chromium negative cases.

chromium negative, concomitant presence of the allergens previously indicated was higher in the first group for every allergen studied ( $p < 0.001$ ) as shown in Table 5.

Seventy eight positive patch tests to chromium (66.10%) were considered to have current or past clinical relevance, with more than one possible source of the exposure in 55 cases. Contact with leather alone or concomitantly with other products, observed in 30 cases (25.42%) was the most frequent association.

Twenty-nine of the 118 (24.58%) cases were workers from the

building industry and 16 (55.17%) reported regular exposure to cement and/or work gloves and/or shoes, 5 of whom only with contact with cement.

When distributing the sources of exposure by gender, we found that contact with cement or cement associated with leather (mostly work gloves) was more frequent in men. On the other hand, women were more frequently exposed to leather or leather concurrently with jewelry, or the latter as the single explainable source of exposure (Table 6).

**Table 6** - Possible sources of exposure to chromium according to gender.

	Total (%)	Men (%)	Women (%)	p-value*
Unknown	53 (44.92%)	23 (42.59%)	30 (46.88%)	-
Cement	8 (6.78%)	8 (14.81%)	0 (0%)	<0.001
Leather	18 (15.25%)	5 (9.26%)	13 (20.31%)	<0.001
Jewelry	7 (5.93%)	1 (1.85%)	6 (9.38%)	<0.001
Cement & Leather	6 (5.08%)	6 (11.11%)	0 (0%)	<0.001
Jewelry & Leather	6 (5.08%)	1 (1.85%)	5 (7.81%)	<0.001
Other Causes	20 (16.95%)	10 (18.51%)	10 (15.63%)	-
<b>Total studied</b>	<b>118 (100%)</b>	<b>54 (100%)</b>	<b>64 (100%)</b>	

\* X<sup>2</sup> test comparing chromium exposure sources between men and women.

## DISCUSSION

The results show a high prevalence (3.60%) of chromium positive patch tests among the population studied for cutaneous hypersensitivity, with a high percentage of current or past relevant suggesting this metal allergen is still a frequent cause of ACD in Portugal, both in the occupational and non-occupational setting. In this population the percentage of positive patch tests is high but in agreement with a previous study performed in the general population without dermatitis in the same region of Portugal in 2010 which showed a prevalence of positive PT to chromium of 1.3%, also with a higher prevalence in males (1.7%) compared to females (1.0%).<sup>19</sup>

A previous study from the same center performed between 1992 and 2011 in the population with cutaneous hypersensitivity found a higher prevalence of chromium allergy - 7%.<sup>12</sup> Although there is a 3-year overlap of data, the lower prevalence of chromium allergy is notorious in our study, especially when considering only the last seven years of the present study that do not overlap with the previous one (3.17% vs 7.0%  $p < 0.001$ ). Nevertheless, after 2010 we found no decreasing tendency in sensitization to chromium. In this period, the directive on the regulation of chromium in cement was already in force since 2005 (DIRECTIVE 2003/53/EC)<sup>17</sup> and the legislation on leather, as we know it today, occurred in 2015 (Commission Regulation (UE) No 301/2014).<sup>17</sup> As expected, the implementation of these measures will have an effect on sensitization only after several years as sensitization will be life-long and, consequently, patch tests will remain positive. Although some of our cases may have had a past relevant many of the workers from the building industry still have more hand dermatitis and aggravation in the occupational setting, namely with cement exposure. Nevertheless, concomitant use of leather gloves may be a confounding factor, because as the chromium directive on leather was later it may take more time to implement it locally. On the other hand, as our study found that occupational exposure was especially related with men over 60 years old, it may suggest these individuals were sensitized by cement in the past, before the implementation of EU directives.

Historically, allergy to this metal was closely related to exposure to cement in male construction workers, but studies from different European countries reveal that this allergy is becoming a consumer problem related to the use of skin products and leather and affecting more the female population,<sup>7</sup> as shown in our studies were positive PT to chromium in females occur mostly outside the occupational setting.

From the occupational point of view, the main sources of exposure are related to the handling of products containing chromium and with individual equipment previously treated with this metal. We are then referring mainly to occupations that manipulate metals, use products such as cement or use individual equipment such as gloves or leather shoes. A frequent co-sensitization to rubber allergens from gloves and shoes, as shown in Table 4, corroborates this hypothesis.

Interestingly, we found only 23.73% of workers from the building industry, which is in line with other European studies. It should also be noted that another study from our center from 1992 to 2011 had found a percentage of 2.3% (123 cases out of 5250) of ACD related to chromium in the building industry, whereas we found only 0.88% (29 out of 3277), all males and with a medium age of 51.69 years.<sup>12</sup> These findings suggest that the application of European Community directives in the regulation of chromium content in cement has been partially effective. It should be noted in Denmark, where chromium limitation in cement occurred long before 2000, a study from 2002 to 2017 did not record cases of ACD from exposure to cement.<sup>6</sup> Concerning the EU directive on limitation of hexavalent chromium in leather there is still no

evident effect, as we have mostly young females with chromium allergy and relation with non-occupational exposure, namely due to shoes. Moreover, frequent association with cobalt and nickel and relation to jewelry may suggest that this metal may also be present in less expensive jewelry after the EU directive limiting exposure to nickel.<sup>20</sup>

This study has limitations that are related to the retrospective nature and the sample of patients with complaints of dermatitis and who sought medical care. This study may represent an image of the current situation and predispose to a more elaborate study that can answer the doubts and hypotheses raised.

## CONCLUSION

In our study, there has not been a significant decrease in sensitization to chromium over the years, but cases classically related to this allergy (cement in the building industry) are no longer the main cause of the problem. Therefore, this suggests that the focus of the problem is on chromium in leather.

**Conflicts of Interest:** The authors have no conflicts of interest to declare. **Financing support:** This work has not received any contribution, grant or scholarship. **Confidentiality of Data:** The authors declare that they have followed the protocols of their work center on the publication of data from patients. **Protection of Human and Animal Subjects:** The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki). **Provenance and Peer Review:** Not commissioned; externally peer reviewed.

**Conflitos de Interesse:** Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho. **Fontes de Financiamento:** Não existiram fontes externas de financiamento para a realização deste artigo. **Confidencialidade dos Dados:** Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes. **Proteção de Pessoas e Animais:** Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial. **Proveniência e Revisão por Pares:** Não comissionado; revisão externa por pares.

### ORCID

Alexandre Afonso: <https://orcid.org/0000-0002-4841-7599>  
Joana Silva: <https://orcid.org/0000-0002-3925-242X>  
Ricardo Silva: <https://orcid.org/0000-0001-5106-7772>  
Vânia Pacheco: <https://orcid.org/0000-0002-4011-1173>  
Vitor Pinheiro: <https://orcid.org/0000-0002-3357-9157>  
Isabel Antunes: <https://orcid.org/0000-0001-9557-0572>  
M. Gonçalves: <http://orcid.org/0000-0001-6842-1360>

### Corresponding Author: Alexandre Afonso

Address: Praceta, R. Prof. Mota Pinto  
3004-561 Coimbra, Portugal  
E-mail: alexan.afonso@hotmail.com

© Author(s) (or their employer(s)) 2021 SPDV Journal. Re-use permitted under CC BY-NC. No commercial re-use.

© Autor (es) (ou seu (s) empregador (es)) 2021 Revista SPDV. Reutilização permitida de acordo com CC BY-NC. Nenhuma reutilização comercial.

## REFERENCES

1. Pestana C, Gomes R, Pinheiro V, Gouveia M, Antunes I, Gonçalves M. Principais causas de dermatite de contacto alérgica ocupacional: Um estudo de três anos no centro de Portugal. *Acta Med Port.* 2016;29:449-55. doi:10.20344/amp.6605
2. Pinheiro V, Pestana C, Marques F, Pinho A, Antunes I, Gonçalves M. Dermite de Contacto Alérgica em Profissionais de Saúde. *J Port Soc Dermatol Venereol.* 2017;75:49-58.

- doi:10.29021/spdv.75.1.718
3. Johansen JD, Frosch PJ, Lepoittevin JP, editors. *Contact Dermatitis*. Berlin: Springer; 2011.
  4. Bourke J, Coulson I, English J. Guidelines for the management of contact dermatitis: An update. *Br J Dermatol*. 2009;160:946-54. doi:10.1111/j.1365-2133.2009.09106.x
  5. Johansen JD, Aalto-Korte K, Agner T, Andersen KE, Bircher A, Bruze M, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing - Recommendations on best practice. *Contact Dermatitis*. 2015;73:195-221. doi:10.1111/cod.12432
  6. Alinaghi F, Zachariae C, Thyssen JP, Johansen JD. Temporal changes in chromium allergy in Denmark between 2002 and 2017. *Contact Dermatitis*. 2019;80:156-61. doi:10.1111/cod.13181
  7. Bregnbak D, Johansen JD, Jellesen MS, Zachariae C, Menné T, Thyssen JP. Chromium allergy and dermatitis: Prevalence and main findings. *Contact Dermatitis*. 2015;73:261-80. doi:10.1111/cod.12436
  8. Tan S, Nixon R. Allergic contact dermatitis caused by chromium in a mobile phone. *Contact Dermatitis*. 2011;65:246-7. doi:10.1111/j.1600-0536.2011.01955.x
  9. Thyssen JP, Jellesen MS, Møller P, Menné T, Johansen JD. Allergic chromium dermatitis from wearing "chromium-free" footwear. *Contact Dermatitis*. 2014;70:185-7. doi:10.1111/cod.12189
  10. Mowitz M, Zimerson E, Hauksson I, Pontén A. Chromate and amine contact allergies in workers manufacturing precast concrete elements. *Contact Dermatitis*. 2016;75:363-9. doi:10.1111/cod.12661
  11. Wong CC, Gamboni SE, Palmer AM, Nixon RL. Occupational allergic contact dermatitis to chromium from cement: Estimating the size of the problem in Australia. *Australas J Dermatol*. 2015;56:290-3. doi:10.1111/ajd.12238
  12. Teixeira V, Coutinho I, Gonçalves M. Alergia de contacto a metais num período de 20 anos no Centro de Portugal: Implicações das directivas Europeias. *Acta Med Port*. 2014;27:295-303. doi:10.20344/amp.4112
  13. Geier J, Krautheim A, Uter W, Lessmann H, Schnuch A. Occupational contact allergy in the building trade in Germany: Influence of preventive measures and changing exposure. *Int Arch Occup Environ Health*. 2011;84:403-11. doi:10.1007/s00420-010-0581-8
  14. Geier J, Lessmann H, Skudlik C, Ballmer-Weber BK, Weisshaar E, et al. Occupational contact allergy in bricklayers, tile setters etc. Current spectrum of sensitization and recent time trends. *Allergol Sel*. 2017;1:127-40. doi:10.5414/alex01593e
  15. Bensefa-Colas L, Stocks SJ, McNamee R, Faye S, Pontin F, Agius RM, et al. Effectiveness of the European chromium(vi) directive for cement implementation on occupational allergic contact dermatitis occurrence: assessment in France and the U.K. *Br J Dermatol*. 2017;177:873-6. doi:10.1111/bjd.15261
  16. Moretto A. Hexavalent and trivalent chromium in leather: What should be done? *Regul Toxicol Pharmacol*. 2015;73:681-686. doi:10.1016/j.yrtph.2015.09.007
  17. European Commission. Commission Regulation (EU) No 301/2014, amending Annex XVII to Regulation (EC) No 1907/2006 of the European Parliament and of the Council on the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) as regards chromium VI compounds. *Off J Eur Union*. 2014:2014-16.
  18. Ministério das Actividades Económicas e do Trabalho. Decreto-Lei no 72/2005 de 18 de Março de 2005. In: *Diário Da República* No 55-I Série A. Lisboa; 2005:2434-2436.
  19. Schuttelaar MLA, Ofenloch RF, Bruze M, et al. Prevalence of contact allergy to metals in the European general population with a focus on nickel and piercings: The EDEN Fragrance Study. *Contact Dermatitis*. 2018;79:1-9. doi:10.1111/cod.12983
  20. European Commission. European Parliament and Council Directive 94/27/EC of 30 June 1994 amending for the 12th time (\*) Directive 76/769/EEC on the approximation of the laws, regulations and administrative provisions of the Member States relating to restrictions on the market. *Off J Eur Communities*. 1994;1993:4-5.